






Molecular evolution and taxonomy of freshwater and terrestrial Ulvophyceae




Pavel Škaloud, Fabio Rindi, Christian Boedeker, Frederik Leliaert, Michala Klimešová,
Veronica Malavasi, Alena Lukešová

Chlorophyta, Ulvophyceae

- usually macroscopic organisms
- prevalently marine algae








Ulvophyceae – thallus types




Ulvophyceae – various ecological habitats


- Marine, brackish, freshwater, terrestrial, epiphytic or endophytic on algae, endozoic in sponges, epizoic on turtles, ...




Ulvophyceae – phylogeny



Ulvophyceae – phylogeny





a




Fučíková et al. 2014

Turmel et al. 2017



0.01




Goals

- Resolving the higher taxonomy of Ulvophytes
- Improving our knowledge on the diversity and evolution of freshwater and terrestrial Ulvophytes
- Taxonomic revision of polyphyletic genera



Methods


- 64 strains from 11 algal culture collections (ACOI, ASIB, CAUP, CCAC, CCALA, CCCryo, CCAP, CPCC, FACHB, SAG, UTEX) – incl. 12 types
- 36 strains from own sampling
- Molecular analyses (18S rDNA, ITS, rbcL, tufA)
- Morphological investigations (cell dimensions, shape of pyrenoids, pattern of filament branching, shape of colonies)




Higher taxonomy

- Resurrecting the order **Chlorocystidales** to make Ulvales and Ulotrichales monophyletic

b



Main habitat	Morphology
terrestrial	unicellular
freshwater	colonial/multicellular (including siphonous)
marine	




Desmochloris


Higher taxonomy

- Validation of the order **Ignatales**

b




Main habitat



Morphology

- unicellular
- colonial/multicellular (including siphonous)


136




b

Ignatius


137




b



c



d




f

Pseudocharacium


Higher taxonomy

- Ulvales:
describing the family


Pseudopleurococcaceae



Pseudopleurococcus



0.02




Main habitat	Morphology
marine	unicellular
freshwater	filamentous
terrestrial	parenchymatous

Ⓐ authentic strains

Higher taxonomy

- **Ulotrichales:** describing the families

Hazeniaceae Planophilaceae Trichosarcinaceae Neocloniaceae Binucleariaceae



Trichosarcina

Problem of types

- Taxonomic position of **Ulothrix**




- Uronema:**

- *Ulothrix acuminata* FACHB 1747
- *Ulothrix sp.* FACHB 781
- *Ulothrix sp.* CPCC 546
- *Ulothrix belkae* SAG 34.86
- *Ulothrix acuminata* UTEX 1178 T






Ulothrix



Uronema

Problem of types

- Taxonomic position of **Ulothrix**
- *Ulothrix tenuissima*
Kützing 1833
- Timavo, Italy



Misidentifications and “molecular” identifications

Binuclearia sp. CCAC 4824

- *Microspora*
- *Klebsormidium*
- *Klebsormidium*
- *Klebsormidium*
- *Klebsormidium*
- *Geminella*
- ...

Binuclearia sp. SAG 39.91

Ulothrix cf. *variabilis* CCCryo 149, 150

Ulothrix sp. CPCC 509, 568

Ulothrix crenulata SAG 35.86


Ulothrix sp. SAG 90.91

...




Misidentifications and "molecular" identifications

- Sluiman (1991): *Gloeotilopsis planctonica* – SAG 29.93
- Friedl (1996): new combinations for *Rhexinema* and *Pleurastrum* species




Misidentifications and "molecular" identifications

- Sluiman (1991): *Gloeotilopsis planctonica* – SAG 29.93
- Friedl (1996): new combinations for *Rhexinema* and *Pleurastrum* species




Misidentifications and “molecular” identifications

- Sluiman (1991): *Gloeotilopsis planctonica* – SAG 29.93
- Friedl (1996): new combinations for *Rhexinema* and *Pleurastrum* species






Misidentifications and “molecular” identifications

- Sluiman (1991): *Gloeotilopsis planctonica* – SAG 29.93
 - Friedl (1996): new combinations for *Rhexinema* and *Pleurastrum* species





Misidentifications and "molecular" identifications

- Friedl (1995): *Pleurastrum insigne* SAG 30.93 in Chlorophyceae
- Kawasaki et al. (2015): SAG 30.93 was overgrown by *Chlorococcum* already in 1994




Misidentifications and “molecular” identifications

- Friedl (1995): *Pleurastrum insigne* SAG 30.93 in Chlorophyceae
- Kawasaki et al. (2015): SAG 30.93 was overgrown by *Chlorococcum* already in 1994




Considerable cryptic diversity

- Pseudendoclonium* morphotype in at least 4 independent lineages




Ecological consequences

- numerous transitions between marine and freshwater habitats
- terrestrial taxa evolved from freshwater relatives
- broad ecological range of *Pseudendoclonium*



Freshwater Flora of Central Europe

- to be released early in 2018



Thank you for your attention!

