

Phyllosiphon duini sp. nov. (Trebouxiophyceae, Chlorophyta), a Species Isolated from a Corticolous Phototrophic Biofilm

Author(s): Kateřina Procházková, Yvonne Némcová & Jiří Neustupa Source: Cryptogamie, Algologie, 39(1):23-34. Published By: Association des Amis des Cryptogames <u>https://doi.org/10.7872/crya/v39.iss1.2018.23</u> URL: <u>http://www.bioone.org/doi/full/10.7872/crya/v39.iss1.2018.23</u>

BioOne (www.bioone.org) is a nonprofit, online aggregation of core research in the biological, ecological, and environmental sciences. BioOne provides a sustainable online platform for over 170 journals and books published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Web site, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/page/terms_of_use.

Usage of BioOne content is strictly limited to personal, educational, and noncommercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Cryptogamie, Algologie, 2018, 39 (1): 23-34 © 2018 Adac. Tous droits réservés

Phyllosiphon duini sp. nov. (Trebouxiophyceae, Chlorophyta), a species isolated from a corticolous phototrophic biofilm

Kateřina PROCHÁZKOVÁ*, Yvonne NĚMCOVÁ & Jiří NEUSTUPA

Department of Botany, Faculty of Science, Charles University, Benátská 2, CZ-12801, Prague, Czech Republic

Abstract – The Watanabea clade of the Trebouxiophyceae includes unicellular coccoid green microalgae that mostly thrive in subaerial microhabitats. Recently, a number of new genera and species were described on the basis of the DNA sequence data and morphological observations. The peculiar genus Phyllosiphon, which forms a monophyletic clade within the Watanabea clade, is characterized by forming siphonous parasitic stages thriving in the leaves of the Araceae. In addition, several previous studies demonstrated that members of the genus Phyllosiphon also include free-living chlorelloid individuals that occur on various subaerial substrates. A number of *Phyllosiphon* members sampled from European subaerial microhabitats have so far not been taxonomically described, because they were not available in cultures. In this study, we provide a taxonomic description of a new species of the genus *Phyllosiphon*, *P. duini*, isolated from a corticolous biofilm growing on *Quercus pubescens* in a sub-Mediterranean forest stand. The simple chlorelloid morphology of this strain did not unambiguously distinguish *Phyllosiphon duini* from other closely related members of the Watanabea clade. However, phylogenetic analyses based on the 18S rDNA sequences showed that this species clustered in a sister position to a *Phyllosiphon* species previously described from eastern Asia. Similar phylogenetic pattern was also supported by the plastid-encoded rbcL gene sequences of members of the Watanabea clade. Our data demonstrate that the genus *Phyllosiphon* represents a diverse phylogenetic lineage within the subaerial chlorelloid green microalgae of the Watanabea clade.

18S rDNA / green algae / *Phyllosiphon / rbc*L / subaerial algae / Trebouxiophyceae / *Watanabea* clade

INTRODUCTION

Phototrophic corticolous biofilms are typically composed of green microalgae belonging to the division Chlorophyta (Stifterová & Neustupa, 2015). The diversity of these microalgae initially appears low due to their simple, unicellular coccoid morphology. However, examination of their DNA sequence data showed that corticolous green microalgae are considerably phylogenetically diversified (Freystein & Reisser, 2010; Rindi *et al.*, 2010; Kulichová *et al.*, 2014). Most notably, the species and genera belonging to the phylogenetically defined *Watanabea* clade in the class Trebouxiophyceae are especially abundant in corticolous and other subaerial microhabitats, such as rock or soil surfaces (Kulichová *et al.*, 2014; Song *et al.*, 2016). Members of the genus *Chloroidium* have often been frequently encountered in corticolous and epilithic microhabitats (Darienko *et al.*, 2010).

^{*} Author for correspondence: suri.kata@seznam.cz

Chloroidium's sister genus Parachloroidium includes two recently described species that are only known from European corticolous biofilms (Neustupa *et al.*, 2013b; Kulichová et al., 2014). The genera Heterochlorella, Heveochlorella, Kalinella and Mysteriochloris were described from tropical South-East Asian corticolous microhabitats (Zhang et al., 2008; Neustupa et al., 2009; Ma et al., 2013; Song et al., 2016). However, both species of the genus Kalinella were also encountered in sub-Mediterranean corticolous microhabitats in Europe (Kulichová et al., 2014). In addition, genotypes belonging to the genus *Heveochlorella* were also reported as frequently occurring folicolous lichen photobionts in tropical habitats of Florida, USA (Sanders et al., 2016). The genera Watanabea and Viridiella, two deep lineages of the Watanabea clade, as well as the genus Polulichloris, and one species of the genus Desertella, have been found in biofilms growing on soil (Albertano et al., 1991; Fučíková et al., 2014; Song et al., 2015). These chlorelloid microalgae all exhibit the relatively small dimensions of cells that reproduce exclusively asexually by unequally sized autospores. Apart from the above mentioned epiphytic, epilithic and edaphic taxa, the *Watanabea* clade also contains a peculiar parasitic genus, *Phyllosiphon*, which is characterized by the presence of parasitic stages that exhibit a siphonous habit (Aboal & Werner, 2011). Formation of these parasitic stages does not seem to be related to sexual process. Parasitic populations of *Phyllosiphon* form branched filaments in the intracellular matrix of the leaf parenchyma of species in the Araceae family (Procházková et al., 2015; 2016). Previously, the only known DNA sequences from the parasitic members of the *Phyllosiphon* were acquired from infected leaves of subtropical and temperate Arisarum vulgare and Arum italicum (Aboal & Werner, 2011; Procházková et al., 2015; 2016). Interestingly, the life cycle of these parasitic species of the genus Phyllosiphon also includes chlorelloid individuals thriving in corticolous biofilms as free-living algae (Procházková et al., 2016). Recently, an additional *Phyllosiphon* species, *P. coccidium*, was described from a tree bark biofilm in China (Song et al., 2016). Moreover, several additional *Phyllosiphon* genotypes that are closely related to the parasitic and free-living species of *Phyllosiphon* were also reported from various European corticolous and epilithic biofilms (Cutler et al., 2013; Hallmann et al., 2013; Procházková et al., 2015). These genotypes may belong to additional undescribed *Phyllosiphon* species that form their parasitic stages in other species or genera of Araceae. However, they cannot be formally described until the strains are isolated into cultures.

In this study, we investigated a single *Phyllosiphon* strain that was isolated from a corticolous biofilm in a single sub-Mediterranean forest habitat. We characterized the morphology and ultrastructure using light and transmission electron microscopy. We also obtained 18S rDNA and *rbcL* sequences to characterize the phylogenetic position of the strain. The results indicate that it represents a previously unknown lineage of the genus *Phyllosiphon*, which we are describing as a new species, *Phyllosiphon duini*.

MATERIALS AND METHODS

Locality, sampling and cultivation

The novel strain was isolated from the phototrophic corticolous biofilm of *Quercus pubescens* growing in a natural forest close to Duino, Italy (GPS coordinates:

 $45^{\circ}46'42''$ N, $13^{\circ}35'37''$ E; altitude: 25 m a.s.l.) in April 2016. The sample was isolated from approximately 1 cm² of tree bark surface taken from northern side of the trunk at a height of 150 cm. The sample was put into a sterile bag and processed carefully to prevent any cross-contamination. The biofilm was scraped using a sterile dissecting needle into 1.5 ml Eppendorf tube and vortex-mixed for 10 seconds with 1.0 ml of sterile liquid Bold's Basal Medium (BBM) and 0.75 mm diameter sterile glass beads. Then, 40 µl of suspension from each Eppendorf tube was placed onto agar-solidified BBM in Petri dishes. After 6 weeks, algal microcolonies with chlorelloid morphology were isolated onto agar-solidified BBM in test tubes.

Light and electron microscopy

Microphotographs of strains were taken under an Olympus BX51 light microscope with a Canon EOS 700D (Canon, Tokyo, Japan). Transmission electron microscopy was performed following the method described by Procházková *et al.* (2015).

DNA extraction, PCR, and sequencing

The genomic DNA of single algal cultures growing in test tubes was isolated following the protocol described by Procházková et al. (2015), or by using the Invisorb Spin Plant Mini Kit (Invitek, Hayward, CA, USA) according to manufacturer's protocol. The extracted solution was diluted to 5-10 ng μ l-1 and used for PCR. The plastid rbcL gene and the nuclear 18S rDNA marker were PCRamplified from the genomic DNA. Consequently, the *rbc*L gene was amplified from the strain belonging to the genus *Phyllosiphon* acquired from the corticolous biofilm using the primers phyllrbcLF (5'-TTCCGTATGACTCCACAACAAGG-3', Procházková et al., 2015) or PRASF1 (5'-ATGGTTCCACAAACAGAAAC-3', Sherwood et al., 2000) and ellaR2 (5'-TCACGACCTTCATTACGAGCTTG-3', Neustupa et al., 2013a). The 18S rDNA sequences were obtained using these primer combinations: (a) phyllos-F1 (5'-CGGAGAGAGGCTTGAGAATCGGCCTT-3') and 1263R (5'-GAACGGCCATGCACCACC-3', Pichrtová et al., 2013): (b) NS1 (5'-GTAGTCATATGCTTGTCT-3', Hamby *et al.*, 1988) and phyllos-R (5'-GGCAGCAAGGCGGGCCGCG-3'); (c) phyllos-F3 (5'-GACTAGGGATCGGC GGGCGTTTCTCGAA-3') and 1636-57R (5'-GGTAGGAGCGACGGGCGGTGTG-3', Katana et al., 2001) or 18L (5'-CACCTACGGAAACCTTGTTACGACTT-3', Hamby et al., 1988). The PCR mix was performed using the methods described by Procházková et al. (2015). The PCR amplification was carried out under the following conditions: initial denaturation at 94°C for 5/4 minutes (rbcL/18S rDNA); 40/35 cycles of denaturation at 95/94°C for 1 minute/45 seconds, annealing at 50/47/54/52/52°C (for primer combination rbcL-203F and rbcL-991R/phyllrbcLF and ellaR2/PRASF1 and ellaR2/NS1 and phyllos-R/phyllos-F1 and 1263R) for 1/1.5 minutes, and elongation at 72°C for 2/2.5 minutes; final extension at 72°C for 10 minutes. The final segment of the 18S rDNA, which was impossible to obtain using the AmpliTaq Gold 360 DNA polymerase (Thermo Fisher Scientific, Waltham, MA, USA), was amplified using the PrimeSTAR Max DNA Polymerase (Takara, Kusatsu, Shiga Prefecture, Japan). The PCR reactions were performed in a total volume of 15 µl containing 7.5 µl of PrimeSTAR Max DNA Polymerase (0.1 U), 5.9 µl of sterile Milli-Q water, and 0.3 µl of each primer (phyllos-F3 and 1636-57R or 18L; 25 pmol). In this case the PCR protocol consisted of 10 seconds at 98°C,

followed by 35 cycles of 15 seconds at 65°C, 30 seconds at 72°C, and a final extension of 5 minutes at 72°C. All PCR products were analysed by electrophoresis on 1% agarose gel, and stained with ethidium bromide. Correctly amplified products were cleaned using the GenElute PCR Clean-Up Kit (Sigma-Aldrich, St. Louis, MO, USA) according to the manufacturer's protocol. The purified PCR products were sequenced with the amplification primers at Macrogen in Amsterdam, Netherlands, or at Biocev in Vestec, Czech Republic. Sequencing reads were assembled and edited using SeqAssem 09/2004 (Hepperle, 2004). The sequences of the newly described species *Phyllosiphon duini* are available in the GenBank database under the accession numbers KY977525 and KY977526.

Phylogenetic analyses

The newly determined 18S rDNA and rbcL sequences were added to alignments published by Procházková et al. (2016), and manually aligned with additional newly published sequences from the GenBank database using MEGA 6 (Tamura et al., 2013). In parallel, we performed the automatically generated 18S rDNA alignment using ClustalW with default settings, integrated in MEGA 6, and subsequently manually edited there. The final 18S rDNA alignments consisted of 1773 and 1778 nucleotides, respectively. The rbcL gene alignment was done manually only, because it did not contain any gaps. The final *rbcL* alignment consisted of 1211 nucleotides. All these alignments are available at https://botany. natur.cuni.cz/neustupa/phyllosiphon duini.html. The appropriate evolutionary models were determined using the Bayesian information criterion (BIC) in MEGA. The BIC selected the GTR+G+I model for the entire 18S rDNA dataset and the 3rd codon position of *rbc*L, the GTR+G model for the 1st codon position of *rbc*L, and the JC+G+I model for the 2nd codon position of *rbc*L. Phylogenetic trees were inferred with Bayesian inference using MrBayes 3.2.2. (Ronquist et al., 2012). Two parallel Markov chain Monte Carlo (MCMC) runs were carried out for 2 million generations, each with one cold and three heated chains. Analyses of the *rbcL* dataset were carried out using a partitioned dataset to assign distinct substitution models to the codon positions. Parameters and trees were sampled every 100th generation for a total of 20000 trees. After visual inspection of log-likelihood values of sampled trees, the initial 5001 trees of each run were discarded, and posterior probabilities of tree bipartitions were calculated on the basis of the consensus of the remaining $30\,000$ (15000 × 2) trees. The maximum likelihood (ML) and weighted maximum parsimony (wMP) analyses for bootstrap supports of individual phylogenetic lineages were calculated using Garli 2.01 (Zwickl, 2006) and PAUP 4.0b10 (Swofford, 2002), respectively. The maximum likelihood (ML) analyses consisted of 100 replicates, using default settings with automatic termination set at 100000 generations, under the unpartitioned 18S rDNA (GTR + G + I model) and partitioned rbcL (GTR + G, JC + G + I, GTR + G + I model for the first, second and third position, respectively) datasets. The wMP bootstrapping (1000 replicates) was performed using heuristic searches, with 1000 random sequence addition replicates, tree bisection and reconnection (TBR) swapping, and random addition of sequences (the number was limited to 10000 for each replicate), with gap characters treated as a fifth character state. The rescaled consistency index was used to assign weight to the characters on a scale of 0-1000. New weights were based on the mean of the fit values for each character over all of the trees in the memory. The phylogenetic trees were graphically adjusted in FigTree 1.3.1 (Rambaut, 2009) and Adobe Illustrator CS3.

RESULTS

Phylogenetic analyses

The determined 18S rDNA sequence from the species described below as *Phyllosiphon duini* comprised 2140 bp, including one intron (424 bp), which was identified as the intron group I based on literature data, was not shared with any known member of the *Phyllosiphon* clade. The entire intron was excluded from the alignment. Phylogenetic analyses based on the manually edited alignment of 18S rDNA sequences of the major trebouxiophycean clades showed that our strain clustered within the monophyletic Phyllosiphon clade of the Watanabea clade in Trebouxiophyceae (Fig. 14) with high statistical support (1.00 BPP /95 ML bootstrap support/100 wMP bootstrap support). Phyllosiphon duini formed a moderately supported lineage (0.94/65/53) together with *Phyllosiphon coccidium* (KT950842) and "Watanabea" sp. (KU320168). Phyllosiphon duini differed from these taxa by 27 and 29 changes, respectively, out of 1773 nucleotide positions of the 18S rDNA alignment. The similar results were given by phylogenic analyses of automatically generated alignment of 18S rDNA sequences. However, a sister position of our strain to a clade comprising *Phyllosiphon coccidium* (KT950842) and "Watanabea" sp. (KU320168) was inferred with Bayesian phylogenetic inference only with a low statistical support (0.56 BPP). *Phyllosiphon duini* differed from these taxa by 27 and 28 changes, respectively, out of 1778 nucleotide positions of the automated 18S rDNA alignment.

The analyses of the *rbcL* gene sequence (Fig. 15) suggested that *Phyllosiphon duini* clustered in a sister position with *Phyllosiphon* sp. *k17* (KR154336) with high statistical support (1.00/100/100). This lineage clustered in a sister position with *Phyllosiphon coccidium* (KT950844) with moderate support (0.78/-/66). *Phyllosiphon duini* differed from *Phyllosiphon* sp. *k17* and *Phyllosiphon coccidium* by 2 and 39 substitutions out of 1211 nucleotide positions of the *rbcL* alignment. This lineage was part of the highly supported (1.00/100/100) *Phyllosiphon* clade of the *Watanabea* clade in Trebouxiophyceae, which also included the *Phyllosiphon ari* lineage (KU640391, KU640392), and a lineage comprising *Phyllosiphon arisari* (KR154334) and *Phyllosiphon* sp. (KR154335).

Morphology and ultrastructure

Phyllosiphon duini strain CAUP H8804 formed elliptical cells, 5.5-9.5 μ m in diameter. The cells reproduced by 2 to 8 autospores (Figs 4-9). Mostly, there was a single large autospore and 1, 3, 5 or 7 smaller autospores produced within a single sporangium (Figs 4-9). Autosporangia were typically 6.9-12 μ m in diameter, however, sporangia with diameter larger than 16 μ m were also rarely, but occasionally, observed. The cells contained a single parietal chloroplast (Figs 10, 12-13), and occasionally divided into two or three lobes both in mature vegetative cells and autospores (Figs 1-4). The chloroplast contained electron-dense plastoglobuli (Fig. 1) and small elliptical pyrenoids that were covered by the starch envelope (Figs 2b). Cells also contained extraplastidial oil droplets (Fig. 10). The cell wall was smooth and thin, composed of two layers. An outermost layer had fibrillar character (Figs 10-11).

Phyllosiphon duini K. Procházková, Y. Němcová & J. Neustupa sp. nov. Figs 1-14

Description: Vegetative cells solitary, uninucleate. Cells elliptical, 5.5-9.5 μ m in diameter. Single parietal chloroplast containing both starch grains and pyrenoids, divided into two or three lobes. Asexual reproduction via 2-8 autospores, 6.9-12(-16) μ m in diameter. Sexual reproduction not observed. Differs from all the other species of the *Watanabea*-clade of Trebouxiophyceae in "GCC" triplet codon of alanine at the position 163 to 165 of the single copy chloroplast-encoded *rbcL* gene sequence, and in a presence of the unique intron within 18S rDNA.

Holotype specimen: Strain CAUP C-H8804, based on strain 4A-2 obtained from the holotype, has been cryopreserved in the Culture Collection of Algae of Charles University Prague (CAUP) (http://botany.natur.cuni.cz/algo/caup.html). The strain has also been deposited in CAUP as an active culture, CAUP H8804.



Figs 1-9. Morphology of *Phyllosiphon duini* sp. nov. **1-3.** Vegetative cells. **4.** Two-celled autosporangium. **5-6.** Four-celled autosporangium. **7.** Six-celled autosporangium. **8.** Eight-celled autosporangium and autospores. **9.** Autosporangium, autospore and vegetative cell. Scale bars: 5 μm.

Type locality: Subaerial biofilm on the bark of a *Quercus pubescens* growing in Duino, Italy (GPS coordinates: 45°46′42″N, 13°35′37″E; altitude: 25 m a.s.l.) in April 2016.

Distribution: North-east Italy; species was only found at the type locality.

Etymology: The species name is derived from Duino, a small town close to the type locality.



Figs 10-13. Ultrastructure of *Phyllosiphon duini* sp. nov. **10.** Autospore with a single nucleus, mitochondria and chloroplast. **11.** Ongoing autosporogenesis; note a single autospore that is already separated by cleavage furrow (the right side of the cell); the pyrenoids are covered by the starch envelope; the autosporangium is partly wrapped in a mother cell wall. **12.** Autosporangium with mature autospores inside; autospores have a single nucleus, plastid containing starch grains, and extraplastidial oil droplets. **13.** Autosporangium with five visible autospores. Note the fibrillar nature of the outer layer of the cell wall. Abbreviations: c, chloroplast; cf, cleavage furrow; cw, cell wall; m, mitochondria; mcw, mother cell wall; n, nucleus; o, oil droplet; py, pyrenoid; s, starch grain. Scale bars: 0.5 μ m (10), 2 μ m (11), 1 μ m (12, 13).

K. Procházková et al.



Fig. 14. Bayesian tree of Trebouxiophyceae based on the 18S rDNA dataset. Numbers at nodes indicate statistical support (BPP > 0.95/ML > 50%/MP > 50%). Thick branches represent nodes receiving the highest BPP support (1.00). The sequence newly acquired in this study are shown in bold. The scale bar shows the estimated number of substitutions per site.



Fig. 15. Bayesian tree of Trebouxiophyceae based on the *rbcL* dataset. Numbers at nodes indicate statistical support (BPP > 0.95/ML > 50%/MP > 50%). Thick branches represent nodes receiving the highest BPP support (1.00). The sequence newly acquired in this study are shown in bold. The scale bar shows the estimated number of substitutions per site.

DISCUSSION

The genus *Phyllosiphon* is a little known member of the trebouxiophycean *Watanabea* clade with a parasitic lifestyle that infects the leaves of various genera of the Araceae (Aboal & Werner, 2011). However, several recent phylogenetically based studies have demonstrated that the members of the *Phyllosiphon* genus also occur as free-living algae on various subaerial substrates throughout Europe and Asia (Cutler *et al.*, 2013; Hallmann *et al.*, 2013; Procházková *et al.*, 2015; 2016, Song *et al.*, 2016). In this paper, by describing a new *Phyllosiphon* species, *Phyllosiphon duini*, which was found in a corticolous biofilm, we are adding a new piece to the puzzle of understanding the diversity of this clade.

The phylogenetic analyses of the 18S rDNA sequences indicated a sister relationship of the newly described species *P. duini* with a lineage containing P. coccidium, a coccoid free-living microalga recently described from a corticolous biofilm in China, and "Watanabea" sp. isolated from a rock surface in China (Song et al., 2016). The rbcL analysis showed that P. duini formed a well-supported lineage with Phyllosiphon sp. k17 (Procházková et al., 2015), which clustered in a sister position to P coccidium (Song et al., 2016). Two isolates of P. duini 4A-2 and k17 were acquired from the same location in Duino, but on different occasions (P. duini was obtained four years later), and are considered as representatives of the same species. Thus, our newly described species P. duini is, according to the 18S rDNA and *rbcL* analyses, closely related to *Phyllosiphon* species that also thrive as freeliving algae in various subaerial microhabitats. Conversely, both known parasitic *Phyllosiphon* species, *P. ari* and *P. arisari*, are more distantly related to *P. duini* and form their own lineages within the genus. Thus, despite a low number of independent observations of *Phyllosiphon* taxa in natural habitats, it is possible that the genus may be phylogenetically differentiated into lineages with free-living chlorelloid species and lineages containing taxa with alternating parasitic and free-living stages in their life cycle.

This notion was further supported by the fact that, unlike parasitic species *Phyllosiphon arisari* and *P. ari* that usually form the free-living populations in the biofilms on trees occurring close to the vascular plants that host their parasitic stages (Procházková *et al.*, 2015; 2016), we did not find any taxa of the Araceae near to the type locality, nor in other places around Duino, despite our efforts in the field to locate them. Thus, it is possible that the free-living taxa, such as *P. duini*, *P. coccidium*, or other so far undescribed species belonging to the genus *Phyllosiphon*, may only form the simple coccoid stages similar to other members of the *Watanabea* clade (Neustupa *et al.*, 2013a;b; Fučíková *et al.*, 2014; Song *et al.*, 2016). However, it is also possible that parasitic stages of these *Phyllosiphon* taxa are formed occasionally.

In conclusion, our phylogenetic analyses and microscopic observations of the 18S rDNA and *rbcL* gene sequences indicated that *P. duini* represents a new species of the genus *Phyllosiphon*. Considering the growing body of knowledge regarding the diversity of this genus, which occurs not only in the leaves of aracean plants, but also in subaerial microhabitats, we expect that additional taxa of the *Phyllosiphon* lineage might be discovered in other little-known phototrophic microbial biofilms, such as those from other areas with a subtropical climate, or in poorly explored tropical ecosystems. The probable high diversity of tropical *Phyllosiphon* taxa is indirectly supported by observations of parasitic populations of this alga in various tropical aracean plants that form thalli that are morphologically distinct from those known from the European genera *Arisarum* or *Arum* (Lagerheim, 1892; Tobler, 1917). Additional field surveys are also necessary in subtropical and temperate regions to ascertain the geographical range and habitat distribution of the already described species of this genus.

Acknowledgements. We would like to thank to our laboratory technician, Lenka Flašková, for her helpful advice and care. We also thank the anonymous reviewers for their critique and recommendations that improved the manuscript and Editage for English language editing. This work was supported by a grant from the Czech Science Foundation P506/12/0955 and by the research project no. 394315 from the internal Charles University Grant Agency.

REFERENCES

- ABOAL M. & WERNER O., 2011 Morphology, fine structure, life cycle and phylogenetic analysis of *Phyllosiphon arisari*, a siphonous parasitic green alga. *European journal of phycology* 46: 181-192.
- ALBERTANO P., POLLIO A. & TADDEI R., 1991 Viridiella fridericiana (Chlorococcales, Chlorophyta), a new genus and species isolated from extremely acid environments. Phycologia 30: 346-354.
- CUTLER N.A., OLIVER A.E., VILES H.A., AHMAD S. & WHITELEY A.S., 2013 The characterisation of eukaryotic microbial communities on sandstone buildings in Belfast, UK, using TRFLP and 454 pyrosequencing. *International biodeterioration & biodegradation* 82: 124-133.
- FREYSTEIN K. & REISSER W., 2010 Green biofilms on tree barks: more than just algae. In: Seckbach J. & Grube M. (eds), Symbioses and Stress. Springer, Heidelberg, pp. 559-573.
- FUČÍKOVÁ K., LEWIS P.O. & LEWIS L.A., 2014 Widespread desert affiliation of trebouxiophycean algae (Trebouxiophyceae, Chlorophyta) including discovery of three new desert genera. *Phycological research* 62: 294-305.
- HEPPERLE D., 2004 SeqAssem[©]. A sequence analysis tool, contig assembler and trace data visualization tool for molecular sequences. Available from: http://www.sequentix.de/.
- HALLMANN C., STANNEK L., FRITZLAR D., HAUSE-REITNER D., FRIEDL T. & HOPPERT M., 2013 — Molecular diversity of phototrophic biofilms on building stone. *FEMS Microbial* ecology 84: 355-372.
- HAMBY R.K., SIM L.E., ISSEL L.E. & ZIMMER E.A., 1988 Direct RNA sequencing: optimization of extraction and sequencing techniques for work with higher plants. *Molecular biology reports* 6: 179-197.
- KATANA A., KWIATOWSKI J., SPALIK K., ZAKRYŚ B., SZALACHA E. & SZYMAŃSKA H., 2001
 Phylogenetic position of *Koliella* (Chlorophyta) as inferred from nuclear and chloroplast small subunit rDNA. *Journal of phycology* 37: 443-451.
- KULICHOVÁ J., ŠKALOUD P. & NEUŠTÚPA J., 2014 Molecular diversity of green corticolous microalgae from two submediterranean European localities. *European journal of phycology* 49: 345-355.
- LAGERHEIM G., 1892 Über einige neue Arten der Gattung *Phyllosiphon* Kühn. *Nuova Notarisia* 3:120-124.
- MA S., HUSS V.A.R., TAN D., SUN X., CHEN J., XIE Y. & ZHANG J., 2013 A novel species in the genus *Heveochlorella* (Trebouxiophyceae, Chlorophyta) witnesses the evolution from an epiphytic into an endophytic lifestyle in tree-dwelling green algae. *European journal of phycology* 48: 200-209.
- NEUSTUPA J., NĚMCOVÁ Y., ELIÁŠ M. & ŠKALOUD P., 2009 Kalinella bambusicola gen. et sp. nov. (Trebouxiophyceae, Chlorophyta), a novel coccoid Chlorella-like subaerial alga from Southeast Asia. *Phycological research* 57: 159-169.
- NEUSTUPA J., NĚMCOVÁ Y., VĚSELÁ J., STEINOVÁ J. & ŠKALOUD P., 2013a Leptochlorella corticola gen. et sp. nov. and Kalinella apyrenoidosa sp. nov.: two novel Chlorella-like green microalgae (Trebouxiophyceae, Chlorophyta) from subaerial habitats. International journal of systematic and evolutionary microbiology 63: 377-387.
- NEUSTUPA J., NĚMCOVÁ Y., VESELÁ J., STEINOVÁ J. & ŠKALOUD P., 2013b Parachloroidium gen. nov. (Trebouxiophyceae, Chlorophyta), a novel genus of coccoid green algae from subaerial corticolous biofilms. *Phycologia* 52: 411-421.

- PICHRTOVÁ M., REMIAS D., LEWIS L.A. & HOLZINGER A., 2013 Changes in phenolic compounds and cellular ultrastructure of Arctic and Antarctic strains of Zygnema (Zygnematophyceae, Streptophyta) after exposure to experimentally enhanced UV to PAR ratio. *Microbial ecology* 65: 68-83.
- PROCHÁZKOVÁ K., NĚMCOVÁ Y., KULICHOVÁ J. & NEUSTUPA J., 2015 Morphology and phylogeny of parasitic and free-living members of the genus *Phyllosiphon* (Trebouxiophyceae, Chlorophyta). *Nova Hedwigia* 101: 501-518.
- PROCHÁZKOVÁ K., ŇĚMCOVÁ Y. & NEUSTUPA J., 2016 Phyllosiphon ari sp. nov. (Watanabea clade, Trebouxiophyceae), a new parasitic species isolated from leaves of Arum italicum (Araceae). Phytotaxa 283: 143-154.
- RAMBAUT A., 2009 *FigTree, Tree Figure drawing tool*, version 1.3.1. Available from: http://tree. bio.ed.ac.uk/software/figtree/.
- RINDI F., AALLALI H.A., LAM D.W. & LÓPEZ-BAUTISTA J.M., 2010 An overview of the biodiversity and biogeography of terrestrial green algae. In: Rescigno V. & Maletta S. (eds), Biodiversity Hotspots. New York, Nova Science Publishers, pp. 105-122.
- RONQUIST F., TESLENKO M., VAN DER MARK P., AYERS D.L., DARLING A., HÖHNA S., LARGET B., LIU L., SUCHARD M.A. & HUELSENBECK J.P., 2012 — MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice across a large model space. Systematic biology 61: 539-542.
- SANDERS W.B., PÉREZ-ÖRTEGA S., NELSEN M.P. & LÜCKING R., 2016 Heveochlorella (Trebouxiophyceae): a little-known genus of unicellular green algae outside the Trebouxiales emerges unexpectedly as a major clade of lichen photobionts in foliicolous communities. European journal of phycology 52: 840-853.
- SHERWOOD A.R., GARBARY D.J. & SHEATH R.G., 2000 Assessing the phylogenetic position of the Prasiolales (Chlorophyta) using *rbcL* and 18S rRNA gene sequence data. *Phycologia* 39: 139-146.
- SONG H., HU Y., ZHU H., WANG Q., LIU G., LU Z., 2016 Three novel coccoid green algae within the Watanabea clade (Trebouxiophyceae, Chlorophyta) from China. International Journal of systematic and evolutionary microbiology 66: 5465-5477.
- SONG H.Y., ZHANG Q., LIU G.X. & HU Z.Y., 2015 Polulichloris henanensis gen. et sp. nov. (Trebouxiophyceae, Chlorophyta), a novel subaerial coccoid green alga. Phytotaxa 218: 137-146.
- SWOFFORD D.L., 2002 PAUP*: phylogenetic analysis using parsimony (and other methods). Sinauer Associates, Sunderland, Massachusetts.
- ŠTIFTEROVÁ A. & NEUSTUPA J., 2015 Community structure of corticolous microalgae within a single forest stand: evaluating the effects of bark surface pH and tree species. *Fottea* 15: 113-122.
- TAMURA K., STECHER G., PETERSON D., FILIPSKI A. & KUMAR S., 2013 MEGA6: Molecular Evolutionary Genetics Analysis, version 6.0. *Molecular biology and evolution* 30: 2725-2729.
- TOBLER F., 1917 Ein neues tropisches *Phyllosiphon*, seine Lebensweise und Entwicklung. *Jahrbuecher fuer wissenschaftliche botanik* 58:1-28.
- ZHANG J.M., HUSS V.A.R., SUN X.P., CHANG K.J. & PANG D.B., 2008 Morphology and phylogenetic position of a trebouxiophycean green alga (Chlorophyta) growing on the rubber tree, *Hevea brasiliensis*, with the description of a new genus and species. *European journal* of phycology 43: 185-193.
- ZWICKL D.J., 2006 Genetic algorithm approaches for the phylogenetic analysis of large biological sequence datasets under the maximum likelihood criterion. Ph.D. dissertation, The University of Texas at Austin.