

# Multigene phylogeny of *Synura* (Synurophyceae) and descriptions of four new species based on morphological and DNA evidence

BOK YEON JO<sup>1\*</sup>, JONG IM KIM<sup>1\*</sup>, PAVEL ŠKALOUD<sup>2</sup>, PETER A. SIVER<sup>3</sup> AND WOONGGHI SHIN<sup>4</sup>

<sup>1</sup>Department of Biology, Chungnam National University, Daejeon 305-764, Korea

<sup>2</sup>Department of Botany, Faculty of Science, Charles University, Benátská 2, Prague 2, CZ-12801, Czech Republic

<sup>3</sup>Department of Botany, Connecticut College, New London, Connecticut 06320, USA

<sup>4</sup>Department of Biology, Chungnam National University, Daejeon 305-764, Korea

(Received 22 February 2016; revised 8 May 2016; accepted 15 May 2016)

We used phylogenetic analyses based on multiple gene sequences (partial nr SSU and LSU rDNA, partial pt LSU rDNA, *psaA* and *rbcL*) from 148 strains (including three outgroups) and scale ultrastructure to examine phylogenetic relationships among species of the colonial genera *Synura* and *Tessellaria*. The phylogenetic tree based on the combined dataset was congruent with ultrastructural characteristics of the scales. *Synura* was divided into three major clades, two including species in section *Synura*, and one representing section *Peterseniae*. One clade, consisting of seven strains of *S. uvella* (section *Synura*), diverged at the base of the genus. The second clade consisted of the remaining species belonging to the section *Synura*. The third clade, containing organisms in the section *Peterseniae* and characterized by scales possessing a keel, was monophyletic with strong support values. Based on our findings, *S. uvella* needs to be in a separate section from other spine-bearing species, and we therefore propose new sectional ranks; *Synura*, *Peterseniae*, *Curtispinae* (presence of body scales with slender spines, tubular scales and caudal scales). We further propose four new species based on phylogenetic analyses and unique scale characters: *S. longitubularis* sp. nov., *S. sungminbooi* sp. nov., *S. soroconopea* sp. nov. and *S. lanceolata* sp. nov. Lastly, we propose a new genus name, *Neotessella*, to replace the invalid use of the name *Tessellaria*.

**Key words:** molecular phylogeny, morphology, *Neotessella*, scale, synurophytes, *Synura*, *Tessellaria*, ultrastructure

## INTRODUCTION

The genus *Synura* Ehrenberg (1834) is an important component of the phytoplankton community in numerous freshwater habitats worldwide (Kristiansen & Preisig, 2007). There are approximately 46 species and subspecies of *Synura*, including two fossil taxa, described from Africa (Hansen, 1996), the Arctic region (Bradley, 1964; Asmund, 1968; Kristiansen, 1992), Asia (Takahashi, 1972, 1973; Kim, 1997; Boo *et al.*, 2010), Europe (Cronberg & Kristiansen, 1980; Hartmann & Steinberg, 1989; Siver *et al.*, 2005; Škaloud *et al.*, 2013, 2014), and North and South America (Nicholls & Gerrath, 1985; Siver, 1987, 1988; Siver & Wujek, 1993, 1999; Siver & Vigna, 1997; Vigna & Munari, 2001; Siver & Wolfe, 2005; Siver, 2013). *Synura* species are colonial flagellates characterized by cells that are covered with siliceous scales, two visible and unequal length flagella and two

plastids. Scale morphology and shape are dependent, in part, on their position on the cell surface, and often consists of apical tubular scales, body scales, transition scales and caudal scales. Because apical and caudal scales in some species are characterized by more cylindrical or elongated morphologies, respectively, distinguishing among species is based largely on differences in body scales (Kristiansen & Preisig, 2007).

The systematics of *Synura* were revolutionized once the ultrastructure of scales was resolved with the electron microscope and the new information led to modifications in the classification of *Synura*, from earlier sectional to detailed serial ranks (Petersen & Hansen, 1956, 1958; Fott & Ludvík, 1957; Asmund, 1968; Balonov & Kuzmin, 1974; Péterfi & Momeu, 1977; Takahashi, 1978; Cronberg, 1989; Škaloud *et al.*, 2012, 2013, 2014). The classification scheme of Petersen & Hansen (1956) divided *Synura* into two major groups, section *Peterseniana* and section *Uvellae*. Species in *Peterseniana* were defined as

Corresponding author: Woongghi Shin. E-mail: shinw@cnu.ac.kr

\*These two authors contributed equally to this work.

†Present address: Advanced Bio-resource Research Center, Kyungpook National University, Daegu 702-701, Korea.

having body scales with a central keel with either a blunt or spine-like projection. Within the section, the identification of species and infraspecific taxa is based primarily upon scale dimension, keel shape, size of the base plate holes and the character of struts. Species in *Uvellae* were defined as having body scales with a forward-projecting spine. Within the section, the identification of species and infraspecific taxa is based primarily upon scale dimension, spine shape and dimension, base plate pores, and the pattern of struts and meshwork on the base plate.

Subsequent to the work of Petersen & Hansen (1956), additional subgeneric distinctions were made (Balonov & Kuzmin, 1974; Péterfi & Momeu, 1977; Takahashi, 1978; Cronberg, 1989). Balonov & Kuzmin (1974) modified Petersen & Hansen's system and split the genus into three sections based on the morphological similarities in silica scales: *Lapponicae*, *Peterseniana* and *Synura*. The section *Lapponicae* was established to account for *S. lapponica*, a taxon with oval to circular scales surrounding the whole colony instead of individual cells as is the case for *Synura*. Under the Balonov & Kuzmin (1974) system section *Synura* was equivalent to the section *Uvellae* proposed by Petersen & Hansen (1956). Péterfi & Momeu (1977) accepted the scheme of Balonov & Kuzmin (1974) and also used the name *Synura* for the type section. They further recognized series *Splendidae* within section *Synura*. Recently, Škaloud *et al.* (2013) transferred *S. lapponica* to the genus *Tessellaria*, and proposed a revised classification system that recognizes five sections: *Peterseniae*, *Spinosae*, *Echinulatae*, *Splendidae* and *Uvellae*.

Most molecular phylogenetic studies of *Synura* have focused on cryptic diversity within the *Synura petersenii* complex. The first molecular analyses were performed by Wee *et al.* (2001), who investigated genetic variability in 15 isolates of *S. petersenii* using the nuclear ITS region. Maximum parsimony (MP) analysis revealed the existence of two well-supported clades. Later, Kynčlová *et al.* (2010) extended the molecular dataset by adding a further 21 ITS sequences from newly isolated strains. The maximum likelihood (ML) analysis revealed six different clades within *S. petersenii*. Boo *et al.* (2010) published a seven-protein gene phylogeny of about 100 *S. petersenii* isolates, and confirmed a high degree of cryptic diversity within this species complex. Taxonomic assessment of observed cryptic diversity in the *S. petersenii* complex was first addressed by Škaloud *et al.* (2012), who recognized six lineages as separate species based on scale morphology and DNA sequence data, *S. americana*, *S. conopea*, *S. glabra*, *S. macropora*, *S. petersenii* and *S. truttae*. Most recently, Škaloud *et al.* (2014) described four additional species within the *S. petersenii* complex based on scale morphology and a combined ITS rDNA, *rbcL* and *cox1* dataset.

The main objectives of this study are to (1) determine the phylogenetic relationships among species

within *Synura* based on increased taxon sampling; (2) revise the family designations within the order Synurales; (3) revise the sections within the genus *Synura*; and (4) describe four new species.

## MATERIALS AND METHODS

### Strains and cultures

Collection information and culture accession numbers for 148 strains (including three outgroup species) used in this study are listed in Table S1. Strains were collected with a 20 µm mesh plankton net (Bokyeong Co., Pusan, Korea) from small ponds in Korea. Strains from outside South Korea were obtained from the following algal culture collections: The Provasoli-Guillard National Center for Culture of Marine Algae and Microbiota (NCMA), USA (<http://ncma.bigelow.org/>); Culture Collection of Algae at the University of Göttingen (SAG), Germany (<http://www.epsag.uni-goettingen.de/>); Culture Collection of Algae of Charles University in Prague (CAUP), Czech Republic (Department of Botany, Charles University in Prague). Culture methods followed those previously described (Jo *et al.*, 2011, 2013). We omitted author citations of all *Synura* species in text, but listed full author citations in Table 3.

### Morphological observations and statistical analyses

Cultured *Synura* strains were observed using an Axio Imager A2 microscope (Carl Zeiss Inc., Hallbergmoos, Germany) equipped with differential interference contrast (DIC) optics. Images were captured with an AxioCam HRc (Carl Zeiss Inc., Hallbergmoos, Germany) photomicrographic system. Cellular dimensions were determined by measuring 25–30 cells of each taxon from photographic images.

For field emission SEM, cells were filtered using nylon membrane filters (Whatman Ltd, Maidstone, UK), rinsed in distilled water, fixed in 1% OsO<sub>4</sub>, dehydrated, prepared and viewed according to Jo *et al.* (2011). Voucher specimens were stored at the Chungnam National University (CNUK) herbarium in Daejeon, Korea. For TEM, cells were air dried onto Formvar coated copper grids. The grids were viewed in a JEM 1010 TEM (JEOL Ltd, Tokyo, Japan) at 80 kV. Images were recorded on Kodak EM Film 4489 (Eastman Kodak Co., Rochester, NY, USA) and scanned to digital format using an Epson Perfection V700 Photo scanner (Epson Korea Co. Ltd, Seoul, Korea). Terminology used to describe ultrastructural features of scales follows Škaloud *et al.* (2014). For each newly described species belonging to the section *Peterseniae*, the following characters were measured in 20 randomly selected scales (in *S. sungminbooi*, only 15 well-developed scales were analysed): (1) scale length, (2) scale width, (3) base hole (also referred to as a foramen pore) diameter, (4) average diameter of a keel pore, (5) average diameter of a base-plate pore, (6) keel width and (7) number of struts. The measurements were performed using the program ImageJ 1.45 s (Schneider *et al.*, 2012). Average values of keel and base-plate pore diameters were calculated by measuring areas of approximately 30 randomly selected pores per scale. Measured data were compared with those obtained for 10 closely related *Synura* species by Škaloud *et al.* (2014). Canonical discriminant analysis of measured

data was performed using Statistica 8.0 (StatSoft, Inc., Tulsa, OK, USA). Graphs were generated using the R statistical software (<http://www.r-project.org/>). The position of *S. petersenii* f. *praefracta* in the canonical discriminant ordination space has resolved by measuring the above-mentioned characters in a scale illustrated by Asmund (1968), in her paper describing this taxon.

#### DNA extraction, amplification, sequencing, alignment and phylogenetic analyses

DNA extraction, PCR amplification, PCR product purification, sequencing and sequence alignments were conducted as described in Jo *et al.* (2011). Sequences for the nuclear-encoded SSU and LSU and plastid LSU rRNA genes were aligned by eye using the secondary structure of the rRNA gene sequences of *Mallomonas annulata* (Bradley) Harris (Wuyts *et al.*, 2001) as a guide. Conserved areas of these genes were readily aligned across taxa and were used for phylogenetic analyses, but unalignable nucleotides were excluded from subsequent phylogenetic analyses as well as from pairwise comparisons. The alignment of *rbcL* and *psaA* gene sequences was based on the alignment of inferred amino acid sequences. Only first and second codons of the *rbcL* gene were analysed since the third codon is known as a fast-evolving site (Škaloud *et al.*, 2013).

Phylogenetic analyses were performed using a combined dataset of 9178 characters (nr SSU = 1638, nr LSU = 2548, pt LSU = 2592, pt *psaA* = 1579 and pt *rbcL* = 825) by two different methods: maximum likelihood (ML) and Bayesian inference (BI). The combined dataset included sequences for all 148 strains (Table S1). Although nuclear ITS1 and ITS2 sequences were also determined, these sequences were used as a barcode to help identify species and were used only for phylogenetic analysis of most species within the section *Peterseniae* except for *S. asmundiae*, *S. bjoerkii* and *S. macracantha* (Škaloud *et al.*, 2012, 2014; see Supplementary Fig. S1). The sequences of three chrysophycean species were included as outgroup taxa and to root the tree. Primer regions and ambiguously aligned regions were removed from the alignment prior to phylogenetic analyses. Prior to ML analysis, the best-fit model for the concatenated dataset was estimated by Bayesian Information criterion (BIC) using MODELTEST 3.7 (Posada & Crandall, 1998), and the GTR + I + G model. We used the GTR + I + G nucleotide model as implemented in RAxML v.7.2.8 (Stamatakis, 2006), and used 200 independent tree inferences with the ‘number of run’ option, with default optimized SPR rearrangement and 25 distinct rate categories to identify the best tree. Statistical support for each branch was obtained from 1000 bootstrap replications using the same substitution model and RAxML program settings. Bayesian analyses were performed using MrBayes 3.2 (Ronquist *et al.*, 2012) with a random starting tree and ran for  $2 \times 10^6$  generations, keeping one tree every 1000 generations. Each analysis was performed using the four Metropolis-coupled Markov Chain Monte Carlo (MC<sup>3</sup>), with  $2 \times 10^6$  generations for each chain. Burn-in point was identified graphically by tracking the likelihoods (Tracer v.1.6; <http://tree.bio.ed.ac.uk/software/tracer/>). The first 100 generations were discarded, and the remaining 1901 trees were used to calculate the posterior

probabilities (PP) of each clade. Trees were visualized using FigTree v.1.4.2 (<http://tree.bio.ed.ac.uk/software/figtree/>).

## RESULTS


### Phylogenetic analyses

The molecular data contained 436 new sequences, including 66 of nr SSU, 75 of nr LSU rDNA, 83 of nr ITS, 77 of pt LSU rDNA, 69 of pt *psaA* and 66 of pt *rbcL* data (Table S1), and 296 published sequences, including 76 of nr SSU, 36 of nr LSU rDNA, 27 of nr ITS and 26 of pt LSU rDNA, 55 of pt *psaA* and 76 of pt *rbcL* data (Table S1).

The consensus Bayesian tree was rooted with three species of Chromulinaceae serving as outgroups. The Bayesian and ML analyses recovered trees with almost identical topology (Fig. 1), except for positions of *Mallomonas insignis* Penard and *Synura uvella* isolates. The resulting phylogenetic tree showed that the colonial genus *Synura* was not closely related to *Tessellaria*, but to *Mallomonas*. The genus *Tessellaria* diverged at the base of the synurophyte clade with strong support values (pp = 1.00, ML = 100).

The genus *Synura* divided into three major lineages (clades A, B and C). The section *Synura* divided into two clades (B and C), each forming a monophyletic lineage. Clade A consisted of members of *S. uvella* and formed a strongly supported clade (pp = 1.00, ML = 100). Clade B consisted of the remaining members of the section *Synura* with moderate to strong support (pp = 1.00, ML = 73). Clade B was divided further into B1 and B2. Subclade B1 consisted of *S. splendida*, which diverged at the base of the clade. Subclade B2 consisted of *S. multidentata*, *S. echinulata*, *S. mammillosa*, *S. mollispina*, *S. spinosa*, *S. sphagnicola*, *S. curtispina*, a new species, *S. longitubularis* and several unidentified taxa. The single strain of *S. multidentata* showed a sister relationship to *S. echinulata* and *S. mammillosa* (pp = 1.00, ML = 100). *Synura mammillosa* was not monophyletic, and two strains, S96B5 and S89C3 showed sister relationships to two strains of *S. echinulata*. The single strain of *S. mollispina* was closely related to *S. spinosa* (pp = 1.00, ML = 100). The five strains of *S. spinosa* formed a monophyletic lineage with strongly supported values (pp = 1.00, ML = 100). The five strains of *S. sphagnicola* formed a monophyletic lineage with strong supported values (pp = 1.00, ML = 100). Five strains of *S. curtispina* were closely related to *S. longitubularis*, *Synura* sp. CCMP847, *Synura* sp. 450 and two Korean strains (pp = 1.00, ML = 100). The two strains of *S. longitubularis* formed a monophyletic lineage with well-supported values (pp = 1.00, ML = 100) and sister to two unidentified Korean strains.

The *Peterseniae* (C) clade formed a strongly supported monophyletic clade (pp = 1.00, ML = 100) and



**Fig. 1.** Consensus Bayesian tree of *Synura* species based on a combined nuclear SSU and LSU rDNA, and plastid LSU rDNA, *psaA*, *rbcL* sequences data. Bayesian posterior probability (pp), maximum-likelihood (ML) bootstrap, maximum-parsimony (MP), and distance values (NJ) are shown above or below the branches. The bold branches indicate strongly supported values (pp = 1.00 and ML = 100%). Scale bar, 0.03 substitutions per site.

subdivided into 14 lineages. The single strain of *S. macracantha* diverged at the base of the *Peterseniae* clade, followed by *S. bjoerkii* and *S. asmundiae*. The single strain of *S. bjoerkii* formed sister relationships with *S. asmundiae*, which included two strains (pp = 1.00, ML = 100). *Synura glabra* diverged next to *S. bjoerkii* and *S. asmundiae*, forming a monophyletic group with strong support (pp = 1.00, ML = 100). Members of *S. americana* and *S. petersenii* diverged after *S. glabra*, but the single strain of *S. macropora* was intermixed together with *S. americana* (pp = 1.00, ML = 90), and their relationship was not resolved. The eight strains of *S. petersenii* formed a monophyletic lineage with strong supported values (pp = 1.00, ML = 100) and showed sister relationships with *S. americana* and *S. macropora*. The five strains of *S. truttae* formed a monophyletic lineage with strong support values (pp = 1.00, ML = 99), and showed sister relationships with three strains of *S. heteropora* and four strains of *S. lanceolata* (pp = 1.00, ML = 96). A third newly described species, *S. sungminbooi*, was related to *S. conopea* and a fourth new taxon, *S. soroconopea* (pp = 1.00, ML = 57). The seven strains of *S. sungminbooi* formed a monophyletic lineage with strongly supported values (pp = 1.00, ML = 100). The new species, *S. soroconopea* was closely related to *S. conopea* (pp = 1.00, ML = 75), and formed a monophyletic lineage with strong support values (pp = 1.00, ML = 100).

### Morphological characteristics

The two sections of the genus *Synura* were distinguishable by ultrastructural characteristics of the scales. Section *Synura* was characterized by a hollow forward-projecting spine at the anterior end of the scale; section *Peterseniae* by a keel (hollow ridge) in the middle of the scale. All species included in this study were identified using scale ultrastructural features based on SEM and TEM (Tables 1 and 2). In addition, we described four new species; *S. longitubularis*, *S. sungminbooi*, *S. soroconopea* and *S. lanceolata* based on morphology of body scales (Figs 2–19).

### Taxonomic descriptions

*Synura lanceolata* B.Y. Jo, W. Shin, J.I. Kim & P. Siver sp. nov. (Figs 2–5).

**DESCRIPTION:** Colonies are globular (40–51 µm in diameter). Cells are pyriform or obovate (16–20 × 10–13 µm), entirely covered with lanceolate to ellipsoidal-shaped scales. Body scales are 2.7–3.94 × 1.2–1.7 µm in dimension, with a posterior rim encircling half to two-thirds of the scale perimeter and with small, evenly spaced basal plate pores 20–36 nm in diameter. A round to oblong-shaped base plate hole is 0.24–0.47 µm in diameter, on the anterior end of the scale. Keel

is 1.6–2.7 × 0.4–0.7 µm, marked through the vertical scale axis and elliptical in shape, with a blunt tip often bearing two to three very short teeth. Meshworks along the base of the keel vary from 54 to 82 nm in diameter. Transverse struts are 24–28 in number, extend from the keel to near the edge of the scale perimeter, and the spacing between struts ranges from 0.15–0.34 µm.

**HOLOTYPE SPECIMEN:** Portion of a single gathering of cells on SEM stub CNU056357, deposited at the Herbarium of Chungnam National University, Daejeon, Korea (CNUK). Fig. 5 is a representative scale from the specimen.

**REPRESENTATIVE DNA SEQUENCES:** GenBank accession no. KP268701.

**ISOTYPE SPECIMEN:** Portion of a single gathering of cells on SEM stub CNU056362, deposited at the Herbarium of Chungnam National University, Daejeon, Korea (CNUK).

**STRAIN INFORMATION:** The live culture of the holotype, deposited as Nojeok020610A at the Culture Collection of Algae of Chungnam National University, Daejeon, Korea.

**TYPE LOCALITY:** Nojeok, Chunpo-myeon, Iksan-si, Jeollabuk-do, Korea (35°56'48"N, 127°01'43"E).

**ETYMOLOGY:** The specific epithet '*lanceolata*' refers to the lanceolate shape of the scale.

*Synura longitubularis* B.Y. Jo, W. Shin, J.I. Kim & P. Siver sp. nov. (Figs 6–11).

**DESCRIPTION:** Colonies are globular (40–50 µm in diameter). Cells are pyriform or obovate (22–28 × 8–12 µm), entirely covered with ovate or ellipsoidal scales. Body scales are 2.2–3.9 × 1.6–2.4 µm in dimension with a posterior thickened rim encircling half to two-thirds of the scale perimeter. There are two different-sized base plate pores on the scale except along the peripheral margin. Larger and rimmed pores with a diameter of 100–135 nm are situated at the posterior part of scale. Smaller pores with a diameter of 60–75 nm are found in the anterior part of scale. Hexagonal secondary mesh-work exists in the distal one-third to half of the scale. Each mesh encloses a small pore and struts extend from the meshwork near to anterior edge of the scale. Spine is 0.9–1.6 × 0.3–0.5 µm in dimension, with distal end tapered to an acute point or bearing two to three small teeth on the tip. Caudal scales are spineless, small, slipper-like, 2.0–2.4 × 0.3–0.6 µm. Tubular scales are linear or slightly bent, 5.1–6.3 × 0.2–0.4 µm.


**HOLOTYPE SPECIMEN:** Portion of a single gathering of cells on SEM stub CNU056359, deposited at the Herbarium of Chungnam National University,

**Table 1.** Summary of the major characteristic features that distinguish taxa of the section *Peterseniae*.

Taxon	Cell size ( $\mu\text{m}$ )	Scale size ( $\mu\text{m}$ )	Keel size ( $\mu\text{m}$ )	Base hole diameter ( $\mu\text{m}$ )	Base-pore pore diameter (nm)	Keel pore diameter (nm)	Number of struts	Distance of struts ( $\mu\text{m}$ )	Interconnection of struts by transverse ribs
<sup>a</sup> <i>S. americana</i> Kynčlová & Škaloud	22–28 × 8–12	3.0–4.2 × 1.7–2.3	1.8–2.3 × 0.5–0.7	0.17–0.34	27–63	54–94	21–24	0.27–0.30	very rare
<sup>a</sup> <i>S. borealis</i> Škaloud & Škaloudová	31–42 × 7–12	4.0–5.8 × 1.6–2.6	3.1–3.6 × 0.9–1.2	0.27–0.55	17–26	54–88	28–38	0.27–0.33	Frequent
<sup>a</sup> <i>S. conopea</i> Kynčlová & Škaloud	20–28 × 8–12	3.3–4.1 × 1.4–1.9	1.6–2.8 × 0.5–0.6	0.19–0.44	25–51	70–125	24–30	0.23–0.26	rare
<sup>a</sup> <i>S. glabra</i> Korshikov emend. Kynčlová & Škaloud	19–28 × 10–14	2.4–3.4 × 1.5–2.4	1.3–1.7 × 0.4–0.5	0.14–0.33	29–41	44–100	17–22	0.25–0.29	none
<sup>a</sup> <i>S. heteropora</i> Škaloud, Škaloudová & Procházková	20–25 × 7–11	2.5–3.8 × 1.1–1.9	2.0–2.8 × 0.5–0.7	0.19–0.42	20–31	49–100	22–28	0.18–0.25	frequent
<sup>a</sup> <i>S. hibernica</i> Škaloud & Škaloudová	26–47 × 6–12	3.4–5.6 × 1.2–2.0	3.4–4.1 × 0.7–0.9	0.14–0.38	18–27	49–89	30–47	0.20–0.25	frequent
<b><i>S. lanceolata</i> Jo, Kim, Siver &amp; Shin sp. nov.</b>	<b>16–20 × 10–13</b>	<b>2.7–3.9 × 1.2–1.7</b>	<b>1.6–2.7 × 0.4–0.7</b>	<b>0.24–0.47</b>	<b>20–36</b>	<b>54–82</b>	<b>26–35</b>	<b>0.15–0.34</b>	<b>none</b>
<sup>a</sup> <i>S. laticarina</i> Škaloud & Škaloudová	21–32 × 7–13	3.1–4.3 × 1.6–2.1	2.5–2.9 × 0.7–1.0	0.20–0.33	19–25	52–76	24–32	0.23–0.26	frequent
<sup>a</sup> <i>S. macropora</i> Škaloud & Kynčlová	17–25 × 8–12	2.2–3.4 × 1.5–2.2	1.6–1.8 × 0.4–0.6	0.16–0.39	50–78	69–137	15–22	0.27–0.36	none
<sup>a</sup> <i>S. petersenii</i> Korshikov emend. Škaloud & Kynčlová	20–31 × 8–12	3.6–4.6 × 1.8–2.3	1.7–2.5 × 0.4–0.6	0.24–0.46	18–30	45–82	26–34	0.24–0.28	frequent
<b><i>S. soroconeopea</i> Jo, Kim, Siver &amp; Shin sp. nov.</b>	<b>18–28 × 10–15</b>	<b>2.0–2.9 × 1.0–1.5</b>	<b>1.3–2.3 × 0.5–0.9</b>	<b>0.20–0.53</b>	<b>26–34</b>	<b>65–104</b>	<b>21–26</b>	<b>0.22–0.34</b>	<b>rare</b>
<b><i>S. sungminhoi</i> Jo, Kim, Siver &amp; Shin sp. nov.</b>	<b>20–31 × 10–15</b>	<b>2.5–3.4 × 1.5–1.8</b>	<b>1.3–2.1 × 0.3–0.8</b>	<b>0.20–0.38</b>	<b>15–24</b>	<b>41–120</b>	<b>28–32</b>	<b>0.19–0.41</b>	<b>rare</b>
<sup>a</sup> <i>S. truttae</i> (Siver) Škaloud & Kynčlová	22–31 × 11–13	3.3–3.8 × 1.5–1.8	1.3–2.0 × 0.4–0.5	0.27–0.56	18–25	47–70	27–33	0.19–0.24	frequent

**Table 2.** Summary of the major characteristic features observable with EM used in this study to distinguish between taxa of the section *Curtisipinae*.

Taxon	Cell size ( $\mu\text{m}$ )	Scale size ( $\mu\text{m}$ )	Spine size ( $\mu\text{m}$ )	Spine tip	Secondary structure, diameter of hexagonal meshwork (nm)	Base plate pore diameter (nm)	Posterior rim	Length of tubular scale ( $\mu\text{m}$ )	Other
<i>S. curtispina</i> (Petersen & Hansen) Asmund	20–25 $\times$ 8–12	3.0–4.3 $\times$ 1.3–3.1	1.0–1.4 $\times$ 0.3–0.4	blunt or with 2–3 apical teeth	distal part of scale covered with hexagonal meshwork, 163–200	85–114	upturned rim	4.1–6.1 $\times$ 0.2–0.3 (n = 18)	each hexagonal meshwork including a small pore
<i>S. echinulata</i> Korshikov	11–20 $\times$ 8–10	2.5–3.0 $\times$ 1.6–2.3	up to 1.2	pointed spine	short and radiating struts	58–76	upturned rim	3.8–7.2 $\times$ 0.2–0.3 <sup>a</sup>	distal part of scale with a sinuous pattern
<i>S. korshikovi</i> D. Kapustin & E. S. Gusev	—	4.0–4.9 $\times$ 2.4–3.1	1.5–2.0 $\times$ 0.6–0.8	flat with 2–3 rows of rounded teeth	whole scale covered with hexagonal meshwork, 197–242	60–106	upturned rim encircling 2/3 of scale perimeter	—	each hexagonal meshwork including a small pore
<i>S. longitubularis</i> Jo, Kim, Siver & Shin sp. nov.	22–28 $\times$ 8–12	2.2–3.9 $\times$ 1.6–2.4	0.9–1.6 $\times$ 0.3–0.5	blunt or with 2–3 apical teeth	distal part of scale covered with hexagonal meshwork, 150–240	80–135	upturned rim	5.1–6.9 $\times$ 0.2–0.3 (n = 13)	each hexagonal meshwork including a small pore
<i>S. mammillosa</i> Takahashi	10–14 $\times$ 8–10	2.1–3.6 $\times$ 1.8–2.5	1.0–1.6 $\times$ 0.2–0.4	pointed spine	short and radiating struts	50–70	upturned rim	2.2–2.7 $\times$ 0.2–0.3 (n = 7)	distal part of scale with a sinuous pattern
<i>S. morusimila</i> W. Pang & Q. Wang	23–25 $\times$ 17–18	2.8–3.6 $\times$ 2.1–2.6	1.1–1.8	round with 10–14 teeth	short and radiating struts	43–60	upturned rim	—	stout, cylindrical spine with ribs at its base
<i>S. sphagnicola</i> (Korshikov)	11–15 $\times$ 8–10	2.3–2.8 $\times$ 1.5–2.2	up to 1.6	pointed spine	no further ornamentation	60–75	uninterrupted upturned rim	5.6 $\times$ 0.2 (n=1)	—
<i>S. spinosa</i> Korshikov	19–27 $\times$ 8–12	2.4–5.2 $\times$ 2.0–3.8	0.8–2.1 $\times$ 0.3–0.4	with two small teeth	distal part of scale covered with hexagonal meshwork, 115–173	regularly arranged large and small pores, between 28–92	upturned rim	5.2–9.7 $\times$ 0.2–0.3 (n = 20)	individual hexagonal meshwork does not contain a pore



**Figs 2–11.** Colony and scale morphology of *Synura lanceolata* Nojeok020610A (Figs 2–5) and *Synura longitubularis* Sinsan101709B (Figs 6–11). **Fig. 2.** SEM image of colony forming cells. **Fig. 3.** SEM image showing top surface of body scale with a median keel, transverse strut and posterior rim. **Fig. 4.** SEM image showing bottom surface of body scale with a base plate hole and numerous base plate pores. **Fig. 5.** TEM image of body scale. **Fig. 6.** SEM image showing the colony of *Synura longitubularis* Sinsan101709B. **Fig. 7.** SEM image showing top surface of body scale with secondary layer near the spine with acute end. **Fig. 8.** SEM image showing bottom surface of body scale with smaller pores and larger pores. **Fig. 9.** TEM image of body scale showing hexagonal secondary meshwork. **Fig. 10.** TEM image of caudal scale. **Fig. 11.** TEM image of apical tubular scale. All scale bars = 1 µm.

Daejeon, Korea (CNUK). **Fig. 9** is a representative scale from the specimen.

REPRESENTATIVE DNA SEQUENCES: GenBank accession no. KP268739.

ISOTYPE SPECIMEN: Portion of a single gathering of cells on SEM stub CNU056364, deposited at the Herbarium of Chungnam National University, Daejeon, Korea (CNUK).

STRAIN INFORMATION: The live culture of the holotype, deposited as Sinsan101709B at the Culture Collection of Algae of Chungnam National University, Daejeon, Korea.


TYPE LOCALITY: Sinsan, Dain-myeon, Uiseong-gun, Gyeongsangbuk-do, Korea ( $36^{\circ}25'24''N$ ,  $128^{\circ}22'52''E$ ).

ETYMOLOGY: The specific epithet ‘*longitubularis*’ refers to longer tubular scale than that of a closely related species, *Synura curtispina*.

*Synura sungminbooi* B.Y. Jo, W. Shin, J.I. Kim & P. Siver sp. nov. (Figs 12–15).

DESCRIPTION: Colonies are globular (30–55 µm in diameter). Cells are pyriform or obovate (20–31 × 10–15 µm), entirely covered with ellipsoidal or obovate scales. Body scales are 2.5–3.4 × 1.5–1.8 µm in dimension, with a posterior rim encircling half to two-thirds of the scale perimeter and with small, evenly spaced and finely perforating base plate pores (11–20 nm in diameter). An oval or irregular-shaped base plate hole is 0.32–0.49 µm at the longest axis, on the anterior end of the scale. Keel is 1.3–2.1 × 0.3–0.8 µm in size, marked through the vertical scale axis and subulate or oblong in shape, with an acute tip. Meshworks along the base of the keel vary from 41 to 120 nm in diameter. Transverse struts are 28–32 in number, extending from the keel near to the edge of the scale perimeter. They are usually not interconnected by transverse ribs and distance between struts ranges from 0.19–0.41 µm.

HOLOTYPE SPECIMEN: Portion of a single gathering of cells on SEM stub CNU056356, deposited at the Herbarium of Chungnam National University,



**Figs 12–19.** Colony and scale morphology of *Synura sungminbooi* Geumgok020610A (Figs 12–15) and *Synura soroconopea* Jakeun030207B (Figs 16–19). **Fig. 12.** SEM image showing the colony of *Synura sungminbooi* Geumgok020610A. **Fig. 13.** SEM image showing top surface of body scale with keel, transverse strut and posterior rim. **Fig. 14.** SEM image showing bottom surface of body scale with a big base plate hole and small pores beneath the keel. **Fig. 15.** TEM image of body scale. **Fig. 16.** SEM image showing whole cell of *Synura soroconopea* Jakeun030207B. **Fig. 17.** SEM image showing top surface of body scale with a wide median keel with acute tip and simple strut. **Fig. 18.** SEM image showing bottom surface of body scale with one big hole and a lot of small pores on the plate. **Fig. 19.** TEM image of body scale. All scale bars = 1  $\mu$ m.

Daejeon, Korea (CNUK). **Fig. 15** is a representative specimen.

REPRESENTATIVE DNA SEQUENCES: GenBank accession no. KP268698.

ISOTYPE SPECIMEN: Portion of a single gathering of cells on SEM stub CNU056361, deposited at the Herbarium of Chungnam National University, Daejeon, Korea (CNUK).

STRAIN INFORMATION: The live culture of the holotype, deposited as Geumgok020610A at the Culture Collection of Algae of Chungnam National University, Daejeon, Korea.

TYPE LOCALITY: Geumgok, Duyeo-ri, Yeosan-myeon, Iksan-si, Jeollabuk-do, Korea ( $36^{\circ}04'10''$ N,  $127^{\circ}05'02''$ E).

ETYMOLOGY: The species is named after Prof. Sung Min Boo, who has greatly contributed to phycological taxonomy in Korea.

*Synura soroconopea* B.Y. Jo, W. Shin, J.I. Kim & P. Siver sp. nov. (Figs 16–19).

DESCRIPTION: Colonies are globular (22–48  $\mu$ m). Cells are pyriform or obovate  $18\text{--}28 \times 10\text{--}15$   $\mu$ m, entirely covered with ellipsoidal or obovate scales. Body scales are  $2.0\text{--}2.9 \times 1.0\text{--}1.5$   $\mu$ m in dimension, with a posterior rim encircled half to two-thirds of the scale perimeter and with small sized, evenly spaced basal plate pores (26–34 nm in diameter). A round to oblong-shaped base hole is 0.20–0.53  $\mu$ m in diameter, marked to the anterior end of the scale. Keel is  $1.3\text{--}2.3 \times 0.5\text{--}0.9$   $\mu$ m, marked through vertical axis of the scale and oblongolate or obovate in shape, with an acute tip. Large, evenly spaced keel meshworks are 65–104 nm in diameter at keel base atop the base plate. Transverse struts are 21–26 in number, extending from the keel near to the edge of the scale and lacking interconnection between transverse struts, and spacing between struts, ranges from 0.22–0.34  $\mu$ m.

**HOLOTYPE SPECIMEN:** Portion of a single gathering of cells on SEM stub CNU056355, deposited at the Herbarium of Chungnam National University, Daejeon, Korea (CNUK). **Fig. 19** is a representative scale from the specimen.

**REPRESENTATIVE DNA SEQUENCES:** GenBank accession no. KP268686.

**ISOTYPE SPECIMEN:** Portion of a single gathering of cells on SEM stub CNU056360, deposited at the Herbarium of Chungnam National University, Daejeon, Korea (CNUK).

**STRAIN INFORMATION:** The live culture of the holotype, deposited as CNU01 (Okgeum121810C) at the Culture Collection of Algae of Chungnam National University, Daejeon, Korea.

**TYPE LOCALITY:** Okgeum pond, Jaenam-ri, Yeosan-myeon, Iksan-si, Jeollabuk-do, Korea ( $36^{\circ}24'73''N$ ,  $128^{\circ}25'13''E$ ).

**ETYMOLOGY:** The specific epithet ‘*soroconopea*’ is derived from the species *S. conopea* and Latin *soro-* (=a sister of).

### Morphological analyses

Three newly described species belonging to the section *Peterseniae* (*S. lanceolata*, *S. soroconopea*, *S. sungminbooi*) are evolutionarily young (**Fig. 1**) and morphologically highly similar. To differentiate them from each other and closely related taxa, we morphologically characterized them by measuring several traits associated with the silica scale structure (**Figs 20–23**). Obvious morphological differences were observed in several traits. All three strains produced relatively small silica scales (**Fig. 20**). The scales of *S. soroconopea* and *S. lanceolata* were the narrowest, about  $1.24 \pm 0.12 \mu\text{m}$  and  $1.42 \pm 0.14 \mu\text{m}$  in width, respectively. In addition, *S. soroconopea* could be well recognized by having the shortest scales among all the species analysed, measuring  $2.51 \pm 0.20 \mu\text{m}$  in length. *Synura sungminbooi* was distinct by having exceptionally small base plate pores ( $15.8 \pm 1.6 \text{ nm}$  in diameter) and the largest ratio of keel pore to base plate pore diameter (**Figs 21, 22**). Finally, the newly described species significantly differed by the scale width/keel width ratio (**Fig. 23**). Whereas the scales of *S. sungminbooi* had a very narrow keel compared with the basal plate (width ratio  $3.1 \pm 0.5$ ), *S. soroconopea* is well characterized by an anteriorly widened keel (width ratio  $1.9 \pm 0.2$ ).

The canonical discriminant analysis (CDA) yielded well-defined groupings based on their morphological data. The analysis indicated strongly significant differentiation among the 13 species (Wilks’s  $\lambda = 0.0008$ ;  $P < 0.00001$ ). The forward stepwise analysis indicated


that all tested morphological characters were significant for species recognition ( $P < 0.00001$ ), and selected the base-plate pore diameter as the best discriminating morphological trait (Partial Wilk’s  $\lambda = 0.18$ ). The average correct discrimination of individual scales on the basis of their morphology reached 90%. The correct discrimination level recovered for *S. lanceolata*, *S. soroconopea* and *S. sungminbooi* was 85%, 95% and 73%, respectively. Specifically, three *S. lanceolata* scales were incorrectly classified as *S. truttae* (2) or *S. heteropora* (1), a single *S. soroconopea* scale was classified as *S. lanceolata*, and four *S. sungminbooi* scales were incorrectly classified as *S. truttae* (2) or *S. heteropora* (2). Overall morphological similarity of *S. lanceolata*, *S. sungminbooi*, *S. truttae* and *S. heteropora* based on the suite of morphometric characters is also evident by their close affinity in CDA plots showing the species correlation with canonical axes (**Figs 24, 25**). Indeed, the smallest squared Mahalanobis distances were observed between *S. sungminbooi* – *S. heteropora*, and *S. lanceolata* – *S. truttae* species pairs (3.98 and 5.37, respectively). The first CDA axis was principally correlated with the base-plate pore diameter (correlation coefficient – 0.85), whereas the second and third axes were correlated with the scale length and number of struts, respectively (correlation coefficients –0.81 and 0.58).

## DISCUSSION

### Phylogenetic relationships between families

Our phylogenetic analyses, combining sequence data for five genes from 115 *Synura* strains and two *Tessellaria* strains, confirmed that *Synura lapponica* is a member of *Tessellaria*, and that the genus *Synura* is divided into three major clades. Section *Peterseniae* forms a monophyletic clade, however taxa belonging to section *Synura* as currently described are split into two groups, one containing *S. uvella* and one containing the remaining species of the section. As reported by Škaloud *et al.* (2013), *Tessellaria* (*Synura*) *lapponica* groups with *T. volvocina*, supporting its removal from the family Synuraceae. Other characteristics support this finding. *Tessellaria* species have radially symmetrical scales with a rim that surrounds the entire scale (Goldstein *et al.*, 2005). In addition, *Tessellaria* differs from *Synura* in terms of disposition of the scales around the colony and not around individual cells, the manner in which the colony undergoes asexual fission without mother cell elongation and the formation of compact colonies within a multilayered thick wall (Goldstein *et al.*, 2005). Therefore, both genera are clearly separated based on molecular and morphological characters.

Playfair (1921) first recognized that a new family was required for the genus *Tessellaria* and created the



**Figs 20–25.** Morphometric analysis of siliceous scales. Newly described species *S. lanceolata*, *S. soroconepea* and *S. sungminbooi* are highlighted. **Figs 20–23.** Comparison of eight morphological traits measured in 13 species belonging to the section *Peterseniae*; average values and standard deviations are given. **Fig. 20.** Scatterplot of scale length versus scale width. **Fig. 21.** Scatterplot of base plate pore diameter versus number of struts. **Fig. 22.** Scatterplot of keel pore to base-plate pore diameter ration versus keel width. **Fig. 23.** Scatterplot of scale width to keel width ratio versus base hole diameter. **Figs 24–25.** Canonical discriminant analysis (CDA) of the entire dataset of measured morphological traits, showing the centroids of species displayed in the first and second (**Fig. 24**) and second and third (**Fig. 25**) CDA coordinate system. Asterisk in Figs 24–25 indicate the position of an iconotype of *S. petersenii* f. *praefracta* (Asmund, 1968).

family Tessellariaceae, lacking a description. However, algal taxonomists overlooked this family (Starmach, 1985; Kristiansen, 1986, 2005; Preisig, 1995) and classified *Tessellaria* as a genus in the family Synuraceae (Kristiansen & Preisig, 2007). In addition, the name *Tessellaria* has been found to be invalid. Given our findings, we propose the following three families within the order Synurales Andersen 1987.

Family I. Mallomonadaceae Diesing 1866.

TYPE GENUS: *Mallomonas* Party 1852.

Family II. Synuraceae Lemmermann 1899 emended B.Y. Jo, J.I. Kim, W. Shin, P. Škaloud & P. Siver.

EMENDED DIAGNOSIS: Spherical or elongate colonies. Cells associated by their posterior ends to free-swimming colonies. Each cell covered by two to four discernible types of the silica scales; apical tubular scales, body scales, transition scales and caudal scales;

two flagella being equal or unequal in length; chloroplasts positioned at the periphery of the cell.

TYPE GENUS: *Synura* Ehrenberg 1834.

Family III. Neotessellaceae B.Y. Jo, J.I. Kim, W. Shin, P. Škaloud & P. Siver fam. nov.

DIAGNOSIS: Spherical colony covered by a silica scale case; scales with rims that encircle the perimeter; scales with or without protuberance. Cells united by their tails (or stalks) to free-swimming colonies. Each cell naked; two flagella being equal or unequal in length; chloroplasts positioned at the periphery of the cell.

TYPE GENUS: *Neotessella* B.Y. Jo, J.I. Kim, W. Shin, P. Škaloud & P. Siver nom. nov.

BASIONYM: *Tessella* Playfair 1915 (*Proc. Linn. Soc. N. S.W.*, 40:315–316, Pl. xlv, Figs 6–7).

TYPE SPECIES: *Neotessella volvocina* (Playfair 1915) B. Y. Jo, J.I. Kim, W. Shin, P. Škaloud & P. Siver comb. nov.

BASIONYM: *Tessella volvocina* Playfair 1915 (*Proc. Linn. Soc. N.S.W.*, 40:315–316, Pl. xlv, Figs 6–7).

SYNONYM: *Tessellaria volvocina* Playfair 1918 (*Proc. Linn. Soc. N.S.W.*, 43:508–509, Pl. lvi, Fig. 4).

TYPE: Playfair 1915, Pl. xlv, Figs 6–7 (iconotype).

TYPE LOCALITY: Lagoons at Pott's Hill near Auburn, Australia.

INCLUDED SPECIES: *Neotessella lapponica* (Skuja 1956) B.Y. Jo, J.I. Kim, W. Shin, P. Škaloud & P. Siver comb. nov.

BASIONYM: *Synura lapponica* Skuja 1956 (*Nov. Act. Reg. Soc. Sci. Upsal.* IV, 16: 275–276).

SYNONYM: *Tessellaria lapponica* (Skuja) Škaloud, Kristiansen & Škaloudová 2013 (*Nord. J. Bot.*, 31: 400).

TYPE: Skuja 1956, pp. 275–276, Pl. 47, Figs 10–14, Pl. 48, Figs 1–2p (iconotype).

TYPE LOCALITY: Swampy ponds and lakes around Abisko, Lappland, Sweden.

REMARKS: Playfair (1915) erected the genus *Tessella*, based on *Tessella volvocina*. The genus name was illegitimate due to homonymy with a diatom genus *Tessella* (Ehrenberg 1838). Playfair (1918) renamed the genus, *Tessellaria*. However, the name *Tessellaria* is an illegitimate homonym of a fossil cycadophyte genus *Tessellaria* (*Tessellaria* (Schimper & Mougeot) C.E.I. von Eichwald; see Eichwald 1860). Therefore, since the original genus *Tessella* was validly published, but with an illegitimate name, we propose the genus *Neotessella* nomen novum as the new replacement name.

INCLUDED SPECIES: *Neotessella volvocina* (Playfair 1915) B.Y. Jo, J.I. Kim, W. Shin, P. Škaloud & P. Siver comb. nov.; *Neotessella lapponica* (Skuja 1956) B.Y. Jo, J.I. Kim, W. Shin, P. Škaloud & P. Siver comb. nov.

#### Phylogenetic relationships within *Synura*

Classification within the genus *Synura* is based mainly on ultrastructure of body scales and historically taxa have been divided into two or three infrageneric ranks (Petersen & Hansen, 1956; Balonov & Kuzmin, 1974; Péterfi & Momeu, 1977). A key diagnostic character is the presence of a keel or spine on the base plate of the scale. Péterfi & Momeu (1977), following the infrageneric classification system of Balonov & Kuzmin (1974), recognized three sections: *Lapponica*, *Peterseniae* and *Synura*, the latter section further divided into two series: *Splendidae* and *Synura*. They included *S. splendida* and *S. sphagnicola* in the series *Splendidae*, and the remaining taxa in the series *Synura*. Further, Péterfi & Momeu (1977) noted that species in series *Splendidae* have scales with a long spine and lack additional secondary structures on the base plate. Our results, which indicate that *S. sphagnicola* is more closely related to strains of *S. longitubularis* and *S. curtispina* than to *S. splendida*, do not support this infrageneric concept. More recently, Škaloud *et al.* (2013) proposed five sections corresponding to monophyletic units on their phylogenetic tree based on nuclear SSU rRNA and plastid *rbcL* gene sequences (*Peterseniae*, *Spinosae*, *Echinulatae*, *Splendidae* and *Uvellae*). Although the monophyletic units are the same as in our study, this system needs to be reevaluated to determine which taxonomic level is more reasonable and acceptable for infrageneric rankings. In addition, the section *Uvellae* cannot be accepted according to Article 22 of International Code of Nomenclature for algae, fungi and plants (McNeill *et al.*, 2012; see <http://www.iapt-taxon.org/nomen/main.php>) because its epithet has not repeated the generic name even though the section includes the type of the adopted and legitimate name of the genus. Therefore, the section *Synura* should be established as the corresponding autonym and include the type species, *S. uvella*.

#### Taxonomic revision of subgeneric ranks for *Synura*

With the addition of four new species there are now 49 accepted species and subspecific taxa for *Synura* (Table 3). Based on our findings, we propose that the 49 taxa be divided into the following three sections: *Peterseniae*, *Synura* and *Curtispinae*.

Section I. *Peterseniae* Petersen & Hansen ex Balonov & Kuzmin 1974, *Bot. Jurn. Leningrad.*, 59, 11, p. 1682.

**Table 3.** Check-list of all previously described species of the genus *Synura*. New species are given in bold.

Section	Taxon	Taxonomic status
<i>Lapponica</i>	<i>Synura lapponica</i> Skuja 1956	Basionym (= <i>Tessellaria lapponica</i> (Skuja) Škaloud, Kristiansen & Škaloudova; Škaloud <i>et al.</i> 2013)
<i>Peterseniae</i>	<i>S. adamsii</i> (Smith 1924) Nygaard 1949	Valid
	<i>S. adamsii</i> f. <i>malabarica</i> Philipose 1953	Valid*
	<i>S. americana</i> Kynčlová & Škaloud 2012	Valid (Škaloud <i>et al.</i> 2012)
	<i>S. australiensis</i> Playfair 1915 emend. Croome & Tyler 1985	Valid
	<i>S. borealis</i> Škaloud & Škaloudová 2014	Valid (Škaloud <i>et al.</i> 2014)
	<i>S. caroliniana</i> Whitford 1943	Valid*
	<i>S. conopea</i> Kynčlová & Škaloud 2012	Valid (Škaloud <i>et al.</i> 2012)
	<i>S. elipidosa</i> Skvortsov 1961	Valid*
	<i>S. glabra</i> Korshikov 1929 emend. Kynčlová & Škaloud 2012	Valid (Škaloud <i>et al.</i> 2012)
	<i>S. heteropora</i> Škaloud, Škaloudová & Procházková 2014	Valid (Škaloud <i>et al.</i> 2014)
	<i>S. hibernica</i> Škaloud & Škaloudová 2014	Valid (Škaloud <i>et al.</i> 2014)
	<i>S. intermedia</i> Bioret ex Kufferath in Conrad 1946	Valid*
	<b><i>S. lanceolata</i> B.Y. Jo, W. Shin, J.I. Kim &amp; P. Siver</b>	Newly described in this study
	<i>S. laticarina</i> Škaloud & Škaloudová 2014	Valid (Škaloud <i>et al.</i> 2014)
	<i>S. longisquama</i> Wujek & Elsner 2000	Valid
	<i>S. macracantha</i> (Petersen & Hansen 1958) Asmund 1968	Valid
	<i>S. macropora</i> Škaloud & Kynčlová 2012	Valid (Škaloud <i>et al.</i> 2012)
	<i>S. petersenii</i> Korshikov 1929 emend. Škaloud & Kynčlová 2012	Valid (Škaloud <i>et al.</i> 2012)
	<i>S. petersenii</i> f. <i>asmundiae</i> Cronberg & Kristiansen 1980	Basionym (= <i>S. asmundiae</i> (Cronberg & Kristiansen) Škaloud, Kristiansen & Škaloudova; Škaloud <i>et al.</i> 2013)
	<i>S. petersenii</i> f. <i>bjoerkii</i> Cronberg & Kristiansen 1980	Basionym (= <i>S. bjoerkii</i> (Cronberg & Kristiansen) Škaloud, Kristiansen & Škaloudova; Škaloud <i>et al.</i> 2013)
	<i>S. petersenii</i> f. <i>bonaeensis</i> Vigna 1979	Synonym (= <i>S. petersenii</i> ; Škaloud <i>et al.</i> 2012)
	<i>S. petersenii</i> f. <i>columnata</i> Siver 1988	Valid
	<i>S. petersenii</i> f. <i>glabra</i> (Korshikov 1929) Kristiansen & Preisig 2007	Synonym (= <i>S. glabra</i> Korshikov 1929 emend. Kynčlová & Škaloud; Škaloud <i>et al.</i> 2012)
	<i>S. petersenii</i> f. <i>kufferathii</i> Petersen & Hansen 1958	Synonym (= <i>S. petersenii</i> Korshikov 1929 emend. Škaloud & Kynčlová; Škaloud <i>et al.</i> 2012)
	<i>S. petersenii</i> f. <i>macracantha</i> Petersen & Hansen 1958	Basionym (= <i>S. macracantha</i> (Petersen & Hansen) Asmund; Asmund 1968)
	<i>S. petersenii</i> f. <i>praefracta</i> Asmund 1968	Valid
	<i>S. petersenii</i> f. <i>taymyrensis</i> Kristiansen 1995	Valid
	<i>S. petersenii</i> f. <i>truttae</i> Siver 1987	Synonym (= <i>S. truttae</i> (Siver 1987) Škaloud & Kynčlová; Škaloud <i>et al.</i> 2012)
	<i>S. petersenii</i> var. <i>glabra</i> (Korshikov 1929) Huber-Pestalozzi 1941	Synonym (= <i>S. glabra</i> Korshikov 1929 emend. Kynčlová & Škaloud; Škaloud <i>et al.</i> 2012)
	<b><i>S. soroconopea</i> B.Y. Jo, W. Shin, J.I. Kim &amp; P. Siver</b>	Newly described in this study
	<b><i>S. sungminbooi</i> B.Y. Jo, W. Shin, J.I. Kim &amp; P. Siver</b>	Newly described in this study
	<i>S. truttae</i> (Siver 1987) Škaloud & Kynčlová 2012	Valid (Škaloud <i>et al.</i> 2012)
	<i>S. virescens</i> (Bory de Saint Vincent 1824) Playfair 1921	Valid*
	<i>S. bioretii</i> Huber-Pestalozzi 1941	Valid (probably = <i>S. sphagnicola</i> or <i>S. curtispina</i> : Nicholls & Gerrath 1985)
<i>Curtispinae</i>	<i>S. biseriata</i> Balonov 1976	Valid*
	<i>S. cronbergiae</i> Siver 2013	Valid (fossil species)
	<i>S. conradii</i> Kufferath in Conrad 1946	Synonym (= <i>S. echinulata</i> Korshikov 1929; Petersen & Hansen 1958)
	<i>S. curtispina</i> (Petersen & Hansen 1956) Asmund 1968	Valid
	<i>S. curtispina</i> f. <i>reticulata</i> Asmund 1968	Synonym (= <i>S. curtispina</i> ; Kristiansen & Lind 1995)
	<i>S. danubiensis</i> (Schiller 1929) Starmach 1968	Invalid (= <i>Synuopsis danubiensis</i> ; Wujek & Thompson 2001)
	<i>S. echinulata</i> Korshikov 1929	Valid*
	<i>S. echinulata</i> f. <i>leptorrhhabda</i> Asmund 1968	Synonym (= <i>S. leptorrhhabda</i> ; Nicholls in Nicholls & Gerrath 1985)
	<i>S. falcata</i> Skvortsov 1961	Valid*
	<i>S. favus</i> Bradley 1964	Synonym (= <i>S. curtispina</i> ; Kristiansen & Lind 1995)
	<i>S. globosa</i> (Schiller 1929) Starmach 1968	Valid
	<i>S. granulosa</i> Playfair 1915	Valid*
	<i>S. granulosa</i> var. <i>pusilla</i> Playfair 1915	Valid*
	<i>S. korshikovii</i> Kapustin & Gusev 2015	Valid

(continued)

**Table 3.** Continued.

Section	Taxon	Taxonomic status
	<i>S. leptorrhapha</i> (Asmund 1968) Nicholls in Nicholls & Gerrath 1985	Valid
	<i>S. lohammarii</i> Skuja 1956	Synonym (= <i>S. splendida</i> ; Péterfi 1967)
	<b><i>S. longitubularis</i> B.Y. Jo, W. Shin, J.I. Kim &amp; P. Siver</b>	<b>Newly described in this study</b>
	<i>S. mammillosa</i> Takahashi 1972	Valid
	<i>S. mollispina</i> (Petersen & Hansen 1956) Péterfi & Momeu 1977	Valid
	<i>S. multidentata</i> (Balonov & Kuzmin 1974) Péterfi & Momeu 1977	Valid
	<i>S. nygaardii</i> (Petersen & Hansen 1956) Kristiansen <i>et al.</i> 1997	Valid
	<i>S. punctulosa</i> Balonov 1976	Valid*
	<i>S. reticulata</i> Lemmermann 1904	Synonym (= <i>S. uvella</i> ; Fott & Ludvík 1957)
	<i>S. reticulata</i> var. <i>minor</i> Chodat, Raineri & Drew 1926	Valid*
	<i>S. reticulata</i> var. <i>verrucosa</i> Pascher 1908	Synonym (= <i>S. uvella</i> ; Fott & Ludvík 1957)
	<i>S. sphagnicola</i> (Korshikov 1927) Korshikov 1929	Valid*
	<i>S. spinosa</i> Korshikov 1929	Valid*
	<i>S. spinosa</i> f. <i>curtispina</i> Petersen & Hansen 1956	Synonym (= <i>S. curtispina</i> ; Asmund 1968)
	<i>S. spinosa</i> f. <i>longispina</i> Petersen & Hansen 1956	Valid
	<i>S. spinosa</i> f. <i>mollispina</i> Petersen & Hansen 1956	Basionym (= <i>S. mollispina</i> ; Péterfi & Momeu 1977)
	<i>S. spinosa</i> f. <i>nygaardii</i> Petersen & Hansen 1956	Synonym (= <i>S. nygaardii</i> ; Kristiansen <i>et al.</i> 1997)
	<i>S. spinosa</i> var. <i>striata</i> Cronberg 1989	Valid
	<i>S. splendida</i> Korshikov 1942	Valid*
<i>Synura</i>	<i>S. morusimila</i> Pang & Wang 2013	Valid
	<i>S. recurvata</i> Siver & Wolfe 2005	Valid (fossil species)
	<i>S. uvella</i> Ehrenberg 1834 emend. Korshikov 1929	Valid
	<i>S. uvella</i> f. <i>turfaceae</i> Steinecke 1916	Combined (= <i>S. sphagnicola</i> ; Fott & Ludvík 1957)
	<i>S. uvella</i> f. <i>typica</i> Pascher 1910	Synonym (= <i>S. uvella</i> ; Korshikov 1929)
	<i>S. uvella</i> var. <i>longipes</i> Virieux 1916	Valid
	<i>S. uvella</i> var. <i>punctata</i> Averintsev 1901	Synonym (= <i>S. sphagnicola</i> ; Fott & Ludvík 1957)
	<i>S. uvella</i> var. <i>reticulata</i> (Lemmermann 1904) Pascher 1910	Synonym (= <i>S. uvella</i> ; Fott & Ludvík 1957)
	<i>S. uvella</i> var. <i>tiszaensis</i> Kiss 1978	Invalid
	<i>S. verrucosa</i> Pascher 1913	Synonym (= <i>S. uvella</i> ; Fott & Ludvík 1957)
Uncertain taxa	<i>S. hyalina</i> Skvortsov 1958	Invalid (no species description in the paper and colonial cell without scales)
	<i>S. hyalina</i> var. <i>aculeata</i> Skvortsov 1958	Invalid (no species description in the paper and cell without scales)
	<i>S. hyalina</i> var. <i>rotundata</i> Skvortsov 1958	Invalid (no species description in the paper and cell without scales)
	<i>S. klebsiana</i> (O. Zacharias 1897) Lemmermann 1899	Invalid (= <i>Chrysosphaerella</i> sp.; Kristiansen & Preisig 2007)
	<i>S. microcrepis</i> Nygaard 1978	Synonym (= <i>Chrysodidymus synuroideus</i> ; Kristiansen & Preisig 2007)
	<i>S. rotundata</i> Skvortsov 1958	Invalid (no species description in the paper, but colonial cells with scales)
	<i>S. urogleniformis</i> (Kisselev 1931) Starmach 1968	Invalid (= <i>Synuropsis danubiensis</i> ; Wujek & Thompson 2001)
	<i>S. uvella</i> f. <i>turfosa</i> Steinecke 1922	Invalid (no description)

TYPE SPECIES: *Synura petersenii* Korshikov 1929, *Arch. Protistenkd.* 67, pp. 283, 284, 285, Pl 11, Figs 31–37.

#### Section II. *Synura*.

EMENDED DIAGNOSIS: Body scales with or without a short, stout, conical-shaped spine bearing teeth on the distal end; larger pores present in the base plate, but lacking along the scale perimeter and under the portion of the scale with secondary ornamentation; a thick posterior rim encircling approximately two-thirds of the scale perimeter and supported by distinct struts; tubular and caudal (slipper) scales absent.

TYPE SPECIES: *Synura uvella* Ehrenb. emend. Korshikov 1929, *Arch. Protistenkd.* 67, pp. 279, 280, 281, Pl 11, Figs 31–37.

REMARKS: Key morphological features include the heavy rim subtended by well-developed secondary struts; a stout, conical spine with 1–4 apical teeth and lack of tubular and caudal (slipper) scales. At present this section consists of at least three taxa: *Synura uvella*, *S. morusimila* and *S. recurvata*.

Section III. *Curtispinae* Jo, Kim, Shin, Škaloud & Siver sect. nov.

DIAGNOSIS: Body scales with a more or less slender spine bearing an acute tip or teeth at its end; smaller and larger pores present on the base plate and secondary meshwork; the posterior rim encircling approximately two-thirds to nearly the entire scale perimeter and lacking supporting struts; tubular and caudal scales may be present.

TYPE SPECIES: *Synura curtispina* (Petersen & Hansen, 1956) Asmund 1968, *Hydrobiologia* 31, pp. 506, 507, 508.

REMARKS: The section is a monophyletic lineage, consisting of all members of previous section *Synura* except for three taxa of the section *Synura* including one fossil species.

#### Comparison of the new species with closely related species

The three new species included in section *Peterseniae*, *Synura lanceolata*, *S. soroconopea* and *S. sungminbooi*, have been considered as genetic variants of *S. petersenii* with very low sequence divergences within a 9-gene DNA alignment (Boo *et al.*, 2010). In our phylogeny *S. lanceolata* was closely related to *S. heteropora*, but was morphologically differentiated from this species by having smaller cell and scale size, more struts on the base plate, and lacking transverse ribs between the struts. *Synura petersenii* f. *praefracta* seems to be similar to *S. lanceolata* in terms of a bluntly pointed keel end with a few minute teeth. However, *S. petersenii* f. *praefracta* is characterized by having longer spine of the keel, presence of transverse ribs between struts, and broader scale. Moreover, the ordination analysis of a scale illustrated by Asmund (1968) along with the description of *S. petersenii* f. *praefracta* revealed its slight morphological differentiation from *S. lanceolata*, as well as other analysed *Synura* species (Figs 24, 25). Strains of *S. soroconopea* formed sister relationships with those of *S. conopea*, but the former species is characterized by a smaller scale size, broader keel, smaller keel meshwork diameter and fewer number of struts. Strains of *S. sungminbooi* have scales typical of the section *Peterseniae*, but they have the smallest base plate pores of all known species. In addition, the pores were irregular in shape under TEM. As a member of the section *Curtispinae*, *S. longitubularis* was closely related to *S. curtispina* in the phylogenetic tree, and morphological characters of both species do overlap. However, the new species was differentiated by slightly bigger hexagonal meshworks and longer tubular scales than those of *S. curtispina*. In addition, *S. longitubularis* differed from *S. spinosa* by means of larger

diameter of hexagonal meshworks and wider base plate pores.

#### Cryptic diversity and taxonomic uncertainty within *Synura*

Although our results further clarify the phylogenetic relationships of species within *Synura*, there still remain areas of uncertainty. The topologies of trees obtained by both ML and Bayesian analyses were identical except for relationships among isolates of each species, but differed from those based on datasets containing nuclear SSU rDNA and plastid *rbcL* (Škaloud *et al.*, 2013), or a concatenated ITS rDNA, *rbcL* and *cox1* combination (Škaloud *et al.*, 2014). For example, in clade B of our tree *Synura mollispina* was closely related to *S. spinosa*, but in a tree presented by Škaloud *et al.* (2013) the two species were not related. *Synura macropora* in the section *Peterseniae* did not separate as a monophyletic lineage from *S. americana* based on a dataset consisting of ITS, *psaA*, *rbcL* and *cox1* (see Figure 1, Škaloud *et al.*, 2012). In our ITS tree (see supplement tree) *S. macropora* clearly separated from strains of *S. americana* as a new lineage. Although *S. macropora* is differentiated from *S. americana* based on the sizes of the keel and base plate pores, our phylogenetic analyses support the inclusion of *S. macropora* in the *S. americana* clade. This discrepancy may be due to omitting fast-evolving genes (e.g. ITS, *cox1*) in our concatenated phylogenetic analyses (Fig. 1). Because these sequences are too variable to align for all *Synura* species, they cannot be used to understand phylogenetic relationships among all taxa of the genus *Synura*, but are more useful for understanding relationships between taxa of the section *Peterseniae*.

Recent molecular analyses of the *Synura petersenii* complex based on either nuclear ITS or multigene sequence data show that *S. petersenii* divides into many well-supported clades (Wee *et al.*, 2001; Kynčlová *et al.*, 2010; Boo *et al.*, 2010), and confirms a high degree of cryptic diversity within this species complex. More recently, Škaloud *et al.* (2012, 2014) performed substantial taxonomic studies on cryptic diversity within *S. petersenii*, and recognized 10 cryptic lineages as separate species. Our results are in good agreement with the species concepts of Škaloud *et al.* (2012, 2014), and we additionally include three new species of this complex, based on combined molecular and morphological evidence. In addition, our multigene phylogeny suggests that the sections *Peterseniae* and *Curtispinae* may include more taxa that need to be designated taxonomically, suggesting even further cryptic diversity.

In summary, the phylogenetic tree based on multigene dataset showed that the colonial genera *Synura* and *Neotessella* do not share a sister relationship, but are paraphyletic, and support a separate family distinction for *Neotessella*. The genus *Synura* divided into three

major clades herein designated as *Peterseniae*, *Synura* and *Curtispinae*. *Synura uvella*, one of two modern species within section *Synura*, diverged at the base of the genus, followed by the evolution of the *Peterseniae* and *Curtispinae* clades. Except for *S. macropora*, our multigene data supported all other species as monophyletic lineages, and indicate further cryptic diversity exists within the genus. Lastly, we propose *Neotessella* to replace the invalid use of the name *Tessellaria*.

## ACKNOWLEDGEMENTS

We wish to thank John McNeill, Michael Wynne and Tony Reznicek for discussions with the taxonomic code. The authors would like to thank Dr S.W. Nam for his assistance with electron microscopy.

## DISCLOSURE STATEMENT

No potential conflict of interest was reported by the author(s).

## FUNDING

This research was supported by the Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, Science and Technology (2015R1A2A2A01003192) to W. Shin and grant DEB-1144098 from the U.S. National Science Foundation to P.A. Siver.

## SUPPLEMENTARY INFORMATION

The following supplementary material is accessible via the Supplementary Content tab on the article's online page at <http://dx.doi.org/10.1080/09670262.2016.1201700>

**Supplementary table S1.** Strains of the genus *Synura* used in this study and the GenBank accession numbers for their nr SSU, nr LSU rDNA, ITS, pt LSU rDNA, *psaA* and *rbcL* gene sequences.

**Supplementary fig. S1.** Unrooted Bayesian tree based on 225 nuclear ITS sequences of the *Peterseniae* species using the GTR + I + G evolutionary model [BaseFreq ( $\pi$ A = 0.2842,  $\pi$ C = 0.2301,  $\pi$ G = 0.1956,  $\pi$ T = 0.2902); RateMatrix (2.5444, 5.4851, 7.0150, 0.8953, 5.4851, 1.0000); Gamma shape = 0.6967; Invariant = 0.3744]. Sequences generated in the present study are in red-coloured bold italic.

## AUTHOR CONTRIBUTIONS

B.Y. Jo and J.I. Kim: algal collection, isolation and culture, planning of experiments, molecular analyses and drafting manuscript; P. Škaloud: image and statistical analyses and editing manuscript; P.A. Siver: original concept, drafting and editing manuscript; W. Shin: original concept, algal collection, planning of experiments, drafting and editing manuscript.


## REFERENCES

- Andersen, R.A. (1987). Synurophyceae classis nov., a new class of algae. *American Journal of Botany*, **74**: 337–353.
- Asmund, B. (1968). Studies on Chrysophyceae from some ponds and lakes in Alaska. VI. Occurrence of *Synura* species. *Hydrobiologia*, **31**: 497–515.
- Averintsev, S. (1901). Materialy k poznavaniju fauny prostejsich v okresnostjach Bologovskoj Biologicheskoy stancii. *Trudy Presnovodnoi Biologicheskoi Stantsii Imperatorskago S.-Peterburgskago Obshchestva Estestvoispytatelei*, **1**: 205–238.
- Balonov, I.M. (1976). Rod *Synura* Ehr. (Chrysophyta). Biologija, Ekologija, Sistematička. Akademija Nauk SSSR, Institut Vnutrennjich Vod, *Trudy*, **31**: 61–82.
- Balonov, I.M. & Kuzmin, G.V. (1974). Vidy roda *Synura* Ehrenberg (Chrysophyta) v vodokhranilischchakh Volzhskogo Kaskada. *Botanicheskiy Zhurnal*, **59**: 1675–1686.
- Boo, S.M., Kim, H.S., Shin, W., Boo, G.H., Cho, S.M., Jo, B.Y., Kim, J.-H., Kim, J.H., Yang, E.C., Siver, P.A., Wolfe, A.P., Bhattacharya, D., Andersen, R.A. & Yoon, H.S. (2010). Complex phylogeographic patterns in the freshwater alga *Synura* provide new insights into ubiquity vs. endemism in microbial eukaryotes. *Molecular Ecology*, **19**: 4328–4338.
- Bory de Saint Vincent, J.B.G.M. (1824). Encyclopédie méthodique. Histoire naturelle des zoophytes, ou animaux rayonnés. Veuve Agasse, Paris. 819 pp.
- Bradley, D.E. (1964). A study of the *Mallomonas*, *Synura* and *Chrysosphaerella* of Northern Iceland. *Journal of General Microbiology*, **37**: 321–333.
- Chodat, R., Raineri, R. & Drew, K. (1926). Algues de la région du Grand Saint-Bernard. *Bulletin de la Société Botanique de Genève, série II*, **17**: 202–217.
- Conrad, W. (1946). Notes protistologiques. XXXI. Matériaux pour la morphologie des *Synura* Ehrenberg. *Bulletin du Musée Royal d'Histoire Naturelle de Belgique*, **22**: 1–12.
- Cronberg, G. (1989). Scaled chrysophytes from the Tropics. *Nova Hedwigia*, **95**: 191–232.
- Cronberg, G. & Kristiansen, J. (1980). Synuraceae and other Chrysophyceae from central Småland, Sweden. *Botaniska Notiser*, **133**: 595–618.
- Croome, R.L. & Tyler, P.A. (1985). *Synura australiensis* (Mallomonadaceae, Chrysophyceae), a light and electron microscopical investigation. *Nordic Journal of Botany*, **5**: 399–401.
- Diesing, K.M. (1866). Revision der Prothelminthen. Abtheilung: Mastigophoren. *Sitzungsberichte – Akademie der Wissenschaften in Wien, Mathematisch-naturwissenschaftliche*, **52**: 287–401.
- Ehrenberg, C.G. (1834). Dritter Beitrag zur Erkenntnis großer Organisation in der Richtung des kleinsten Raumes. *Abhandlungen der Königlichen Akademie der Wissenschaften zu Berlin*, **1833**: 145–336.
- Ehrenberg, C.G. (1838). *Die infusionsthierchen als vollkommene Organismen. Ein Blick in das tiefere organische Leben der Natur*. Leopold Voss Verlag, Leipzig. 244 pp.
- Eichwald, E. (1860). *Lethaea Rossica ou Paléontologie de la Russie*. Stuttgart (Schweizerbart). 1654 pp.
- Fott, B. & Ludvík, J. (1957). Die submikroskopische Struktur der Kieselalgen bei *Synura* und ihre Bedeutung für die Taxonomie der Gattung. *Preslia*, **29**: 5–16.
- Goldstein, M., McLachlan, J. & Moore, J. (2005). Morphology and reproduction of *Synura lapponica* (Synurophyceae). *Phycologia*, **44**: 566–571.
- Hansen, P. (1996). Silica-scaled Chrysophyceae and Synurophyceae from Madagascar. *Archiv für Protistenkunde*, **147**: 145–172.
- Hartmann, H. & Steinberg, C. (1989). The occurrence of silica-scaled chrysophytes in some central European lakes and their relation to pH. *Nova Hedwigia*, **95**: 131–158.
- Huber-Pestalozzi, G. (1941). Das Phytoplankton des Süßwassers. 2,1. Chrysophyceen, farblose Flagellaten, Heterokonten. In *Die Binnengewässer* (Thienemann, A., editor), Band XVI. Schweizerbart'sche Verlagsbuchhandlung, Stuttgart. 365 pp.
- Jo, B.Y., Shin, W., Boo, S.M., Kim, H.S. & Siver, P.A. (2011). Studies on ultrastructure and three-gene phylogeny of the genus *Mallomonas* (Synurophyceae). *Journal of Phycology*, **47**: 415–425.
- Jo, B.Y., Shin, W., Kim, H.S., Siver, P.A. & Andersen, R.A. (2013). Phylogeny of the genus *Mallomonas* (Synurophyceae) and

- descriptions of five new species on the basis of morphological evidence. *Phycologia*, **52**: 266–278.
- Kapustin, D. A. & Gusev, E.S. (2015). *Synura korshikovii* sp. nov. (Chrysophyceae, Synurales), a new species from Ukraine. *Phytotaxa*, **232**: 185–190.
- Kim, H.S. (1997). Silica-scaled chrysophytes (Synurophyceae) in several reservoirs, swamps, and a highland pond from Changnyong County, Korea. *Algae*, **12**: 1–10.
- Kiss, I. (1978). Occurrence of *Synura uvella* Ehr. var. *tiszaensis* n. var. in the dead arm of the river Tisza near Lakitelek. *Tisca*, **13**: 49–54.
- Korshikov, A.A. (1927). *Skadovskiella sphagnicola*, a new colonial chrysomonad. *Archiv für Protistenkunde*, **58**: 450–455.
- Korshikov, A. (1929). Studies on the chrysomonads I. *Archive für Protistenkunde*, **67**: 253–290.
- Korshikov, A.A. (1942). On some new or little known flagellates. *Archiv für Protistenkunde*, **95**: 22–44.
- Kristiansen, J. (1986). The ultrastructural bases of chrysophyte systematics and phylogeny. *CRC Critical Reviews in Plant Sciences*, **4**: 149–211.
- Kristiansen, J. (1992). Silica-scaled chrysophytes from West Greenland: Disko Island and the Søndre Strømfjord region. *Nordic Journal of Botany*, **12**: 525–536.
- Kristiansen, J. (1995). Silica-scaled chrysophytes from Lake Thingvallavatn, Iceland. *Archiv für Hydrobiologie, Supplement 111/Algological Studies*, **79**: 67–76.
- Kristiansen, J. (2005). *Golden Algae: A Biology of Chrysophytes*. A. R.G. Gantner Verlag. 167 pp.
- Kristiansen, J. & Lind, J.F. (1995). On the taxonomic relation between *Synura curtispina* and *S. favus* (Synurophyceae). *Nordic Journal of Botany*, **15**: 443–447.
- Kristiansen, J. & Preisig, H.R. (2007). Chrysophyte and Haptophyte Algae, 2: Teil/Part 2: Synurophyceae. In *Süßwasserflora von Mitteleuropa* (Büdel, B., Gärtner, G., Krienitz, L., Preisig, H.R. & Schagerl, M., editors), vol. 1/2. Springer-Verlag, Berlin. 252 pp.
- Kristiansen, J., Düvel, L. & Wegeberg, S. (1997). Silica-scaled chrysophytes from the Taymyr Peninsula, Northern Siberia. *Nova Hedwigia*, **65**: 337–351.
- Kynčlová, A., Škaloud, P. & Škaloudová, M. (2010). Unveiling hidden diversity in the *Synura petersenii* species complex (Synurophyceae, Heterokontophyta). *Nova Hedwigia*, **136**: 283–298.
- Lemmermann, E. (1899). Das Phytoplankton sächsischer Teiche. *Forschungsberichte aus der Biologischen Station zu Plön*, **7**: 96–135.
- Lemmermann, E. (1904). Das Plankton schwedischer Gewässer. *Arkiv för Botanik, Uppsala*, **2**: 1–209.
- McNeill, J., Barrie, F.R., Buck, W.R., Demoulin, V., Greuter, W., Hawksworth, D.L., Herendeen, P.S., Knapp, S., Marhold, K., Prado, J., Prud'homme Van Reine, W.F., Smith, G.F., Wiersema, J.H. & Turland, N.J. (2012). International Code of Nomenclature for algae, fungi, and plants (Melbourne code) adopted by the Eighteenth International Botanical Congress Melbourne, Australia, July 2011. Regnum Vegetabile 154. Koeltz Scientific Books, Germany.
- Nicholls, K.H. & Gerrath, J.F. (1985). The taxonomy of *Synura* (Chrysophyceae) in Ontario with special reference to taste and odour in water supplies. *Canadian Journal of Botany*, **63**: 1482–1493.
- Nygaard, G. (1949). Hydrobiological studies on some Danish ponds and lakes. *Kongelige Danske Videnskabernes Selskab, Biologiske Skrifter*, **7**: 1–293.
- Nygaard, G. (1978). Freshwater phytoplankton from the Narssaq area, South Greenland. *Botanisk Tidsskrift*, **73**: 191–238.
- Pang, W. & Wang, Q. (2013). A new species, *Synura morusimila* sp. nov. (Chrysophyta), from Great Xing'an Mountains, China. *Phytotaxa*, **88**: 55–60.
- Pascher, A. (1908). Ein kleiner Beitrag zur Kenntnis der Chrysomonaden Böhmens. *Lotos*, **57**: 148–154.
- Pascher, A. (1910). Der Grossteich bei Hirschberg in Nord-Böhmen. I. Chrysomonaden. *Monographien und Abhandlungen zur Internationale Revue der gesamten Hydrobiologie und Hydrographie*, **1**: 1–66.
- Pascher, A. (1913). Chrysomonadinae. In *Die Süßwasser-Flora Deutschlands, Oesterreichs und der Schweiz* (Ed. by A. Pascher, A., editor), Vol. 2. G. Fischer, Jena. 7–95 pp.
- Perty, M. (1852). *Zur Kenntnis kleinsten Lebensformen nach Bau, Funktionen, Systematik, mit Specialverzeichniss der in der Schweiz beobachteten*. Bern, Switzerland. 228 pp.
- Péterfi, L.S. (1967). Studies on the Rumanian Chrysophyceae. I. *Nova Hedwigia*, **13**: 117–137.
- Péterfi, L.S. & Momeu, L. (1977). Remarks on the taxonomy of some *Synura* species based on the fine structure of scales. *Studii si Comunicări Muzeul Brukenthal Știinte Naturale*, **21**: 15–23.
- Petersen, J.B. & Hansen, J.B. (1956). On the scales of some *Synura* species. *Biologiske Meddelelser, Kongelige Danske Videnskabernes Selskab*, **23**(2): 3–27.
- Petersen, J.B. & Hansen, J.B. (1958). On the scales of some *Synura* species. II. *Biologiske Meddelelser, Kongelige Danske Videnskabernes Selskab*, **23**(7): 1–13.
- Philipose, M.T. (1953). Contribution to our knowledge of Indian algae. I. Chrysophyceae. *Proceedings of the Indian Academy of Sciences B*, **37**: 331–338.
- Playfair, G.I. (1915). Freshwater algae of the Lismore District: with an appendix on the algal fungi and Schizomycetes. *Proceedings of the Linnean Society of New South Wales*, **40**: 310–362.
- Playfair, G.I. (1918). New and rare freshwater algae. *Proceedings of the Linnean Society of New South Wales*, **43**: 497–543.
- Playfair, G.I. (1921). Australian freshwater flagellates. *Proceedings of the Linnean Society of New South Wales*, **46**: 99–146.
- Posada, D. & Crandall, K.A. (1998). MODELTEST: testing the model of DNA substitution. *Bioinformatics*, **14**: 817–818.
- Preisig, H.R. (1995). A modern concept of chrysophyte classification. In *Chrysophyte Algae: Ecology, Phylogeny and Development* (Sandgren, C.D., Smol, J.P. & Kristiansen, J.), 46–74. Cambridge University Press, Cambridge.
- Ronquist, F., Teslenko, M., van der Mark, P., Ayres, D., Darling, A., Höhna, S., Larget, B., Liu, L., Suchard, M.A. & Huelsenbeck, J.P. (2012). MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice across a large model space. *Systematic Biology*, **61**: 539–542.
- Schiller, J. (1929). Neue Chryso- und Chryptomonaden aus Altwässern der Donau bei Wien. *Archiv für Protistenkunde*, **66**: 436–458.
- Schimper, W.P. & Mougeot, A. (1844). *Monographie des plantes fossiles du grès bigarré de la chaîne des Vosges*. G. Engelmann, Leipzig. pp. 63–65.
- Schneider, C.A., Rasband, W.S. & Eliceiri, K.W. (2012). NIH image to ImageJ: 25 years of image analysis. *Nature Methods*, **9**: 671–675.
- Siver, P.A. (1987). The distribution and variation of *Synura* species (Chrysophyceae) in Connecticut, USA. *Nordic Journal of Botany*, **7**: 107–116.
- Siver, P.A. (1988). A new form of the common chrysophycean alga *Synura petersenii*. *Transactions of the American Microscopy Society*, **107**: 380–385.
- Siver, P.A. (2013). *Synura cronbergiae* sp. nov., a new species described from two Paleogene maar lakes in northern Canada. *Nova Hedwigia*, **97**: 179–187.
- Siver, P.A. & Vigna, M.S. (1997). The distribution of scaled chrysophytes in the delta region of the Paraná River, Argentina. *Nova Hedwigia*, **64**: 421–453.
- Siver, P.A. & Wolfe, A.P. (2005). Scaled Chrysophytes in Middle Eocene lake sediments from northwestern Canada, including descriptions of six new species. In *Proceedings of the Sixth International Chrysophyte Symposium*. pp. 295–308.
- Siver, P.A. & Wujek, D.E. (1993). Scaled Chrysophyceae and Synurophyceae from Florida, USA. IV. The flora of Lower Lake Myakka and Lake Tarpon. *Florida Scientist*, **56**: 109–117.
- Siver, P.A. & Wujek, D.E. (1999). Scaled Chrysophyceae and Synurophyceae from Florida, USA. VI. Observations on the flora from water bodies in the Ocala National Forest. *Nova Hedwigia*, **68**: 75–92.

- Siver, P.A., Voloshko, L.N., Gavrilova, O.V. & Getsen, M.V. (2005). The scaled chrysophyte flora of the Bolshezemelskaya tundra (Russia). *Nova Hedwigia*, **128**: 125–149.
- Škaloud, P., Kynčlová, A., Benada, O., Kofroňová, O. & Škaloudová, M. (2012). Toward a revision of the genus *Synura*, section Petersenianae (Synurophyceae, Heterokontophyta): morphological characterization of six pseudo-cryptic species. *Phycologia*, **51**: 303–329.
- Škaloud, P., Kristiansen, J. & Škaloudová, M. (2013). Developments in the taxonomy of silica-scaled chrysophytes – from morphological and ultrastructural to molecular approaches. *Nordic Journal of Botany*, **31**: 385–402.
- Škaloud, P., Škaloudová, M., Proncházková, A. & Némcová, Y. (2014). Morphological delineation and distribution patterns of four newly described species within the *Synura petersenii* species complex (Chrysophyceae, Stramenopiles). *European Journal of Phycology*, **49**: 213–229.
- Skuja, H. (1956). Taxonomische und biologische Studien über das Phytoplankton schwedischer binnengewässer. *Nova Acta Regiae Societatis Scientiarum Upsaliensis, Series 4*, **16**: 1–404.
- Skvortsov, B.V. (1958). New and rare Flagellatae from Manchuria, Eastern Asia. *Philippine Journal of Science*, **86**: 139–202.
- Skvortsov, B.V. (1961). Harbin Chrysophyta, China boreali-orientalis. *Bulletin of the Herbarium of North-Eastern Forestry Academy*, **3**: 1–70.
- Smith, G.M. (1924). Ecology of the plankton algae in the Palisades Interstate Park, including the relation of control methods to fish culture. *Roosevelt Wild Life Bulletin*, **2**: 94–195.
- Stamatakis, A. (2006). RAxML-VI-HPC: maximum likelihood-based phylogenetic analyses with thousands of taxa and mixed models. *Bioinformatics*, **22**: 2688–2690.
- Starmach, K. (1968). Chrysophyta I. Chrysophyceae-Zlotowiciowce. *Flora Śląskowodna Polski*, **5**: 1–598.
- Starmach, K. (1985). Chrysophyceae und Haptophyceae. In *Süßwasserflora von Mitteleuropa* (Ettl, H., Gerloff, J., Heynig, H. & Mollenhauer, D., editors), Vol. 1, 1–515. Fischer, Stuttgart.
- Steinecke, F. (1916). Die Algen des Zehlaubruches in systematischer und biologischer Hinsicht. *Schriften der königlichen physikalischökonomischen Gesellschaft zu Königsberg*, **56**: 1–138.
- Steinecke, F. (1922). Die Algen des Pakledimmer Hochmoors. *Botanisches Archiv*, **1**: 225–229.
- Takahashi, E. (1967). Studies on genera *Mallomonas*, *Synura* and other plankton in fresh-water with the electron microscope. VI. Morphological and ecological observations on genus *Synura* in ponds and lakes in Yamagata Prefecture. *Bulletin of the Yamagata University Agricultural Science*, **5**(2): 99–118 [199–218].
- Takahashi, E. (1972). Studies on genera *Mallomonas* and *Synura*, and other plankton in freshwater with electron microscope. VIII. On three new species of Chrysophyceae. *Botanical Magazine Tokyo*, **85**: 293–302.
- Takahashi, E. (1973). Studies on genera *Mallomonas* and *Synura*, and other plankton in fresh water with the electron microscope VII. New genus *Spiniferomonas* of the Synuraceae (Chrysophyceae). *Botanical Magazine Tokyo*, **86**: 75–88.
- Takahashi, E. (1978). *Electron Microscopical Studies of the Synuraceae (Chrysophyceae) in Japan; Taxonomy and Ecology*. Tokai University Press, Tokyo.
- Vigna, M.S. (1979). Ultraestructura de las escamas de *Synura petersenii* Korsh. f. *bonaerensis* nov. forma (Ochromonadales). *Boletín de la Sociedad Argentina de Botánica*, **18**: 47–51.
- Vigna, M. S. & Munari, C. (2001). Seasonal occurrence of silica scales chrysophytes in a Buenos Aires lake. *Nova Hedwigia*, **122**: 195–209.
- Virieux, J. (1916). Recherches sur le plancton des lacs du Jura central. *Annales de Biologie Lacustre*, **8**: 1–192.
- Wee, J.L., Fasone, L.D., Sattler, A., Starks, W.W. & Hurley, D.L. (2001). ITS/5.8S DNA sequence variation in 15 isolates of *Synura petersenii* Korshikov (Synurophyceae). *Nova Hedwigia*, **122**: 245–258.
- Whitford, L.A. (1943). The fresh-water algae of North Carolina. *Journal of the Elisha Mitchell Scientific Society*, **59**: 131–170.
- Wujek, D. & Elsner, P. (2000). *Synura longisquama*, sp. nov., a new synurophyte from South Carolina. *Journal of the Elisha Mitchell Scientific Society*, **116**: 97–100.
- Wujek, D. & Thompson, R. (2001). The chrysophyte genera *Synuropsis* Schiller, *Volvochrysis* Schiller, *Synochromonas* Korshikov, *Pseudosynura* Kisselew, *Chrysomoron* Skuja and *Syncrypta* Ehrenberg. *Transactions of the Kansas Academy of Science*, **104**: 79–91.
- Wuyts, J., De Fijk, P., Van de Peer, Y., Winkelmann, T. & De Wachter, R. (2001). The European large subunit ribosomal RNA database. *Nucleic Acid Research*, **29**: 175–177.
- Zacharias, O. (1897). Neue Beiträge Kenntnis des Süßwasserplanktons. *Forschungsberichte aus der Biologischen Station zu Plön*, **5**: 1–9.

ntaxa = 227  
nchar = 557



**Table S1.** Strains of the genus *Synura* used in this study and the GenBank accession numbers for their nr SSU, nr LSU rDNA, ITS, pt LSU rDNA, *psaA* and *rbcL* gene sequences.

Taxon	Strain	Collection site (GPS coordinates )			GenBank accession		
		Nuclear		Plastid	Plastid	Nuclear	
		SSU	LSU	<i>psaA</i>	<i>rbcL</i>	ITS	
<b><i>Mallomonas</i></b>							
<i>M. adamas</i> Harris & Bradley	CCMP1783	Freshwater reservoir, Mount Macedon, Victoria Australia (37°28'37"S, 144°33'36"E)	JQ281515	JQ281508	KM817930	KM818008	JQ281502
<i>M. akrokomas</i> Ruttner in Pascher	Posan012608J	Posanji, Jindo-eup, Jindo-gun, Jeollanam-do, Korea (34°28'62"N, 126°14'32"E),	GU935625	GU935647	KM817932	KM818010	GU935667
<i>M. annulata</i> (Bradley) Harris	Ilwang022407C	Ihwang, Songjuk-dong, Jangan-gu, Suwon-si, Gyeonggi-do, Korea (37°18'00"N, 126°59'47"E)	GU935626	GU935648	KM817938	KM818016	GU935668
<i>M. areolata</i> Nygaard	Hongseong120107D	Hongseong, Hongseong-eup, Hongseong-gun, Chungcheongnam-do, Korea (36°35'40"N, 126°40'28"E)	GU935619	GU935641	KM817939	KM818017	GU935661
<i>M. asmundae</i> (Wujek & Van der Veer) Nicholls	CCMP 1658	Volo Bog Nature Preserve, McHenry County, IL, USA (42°21'05"N, 88°11'04"W)	M87333	AF409122	-	-	AF015585
<i>M. bangladeshica</i> (Takahashi & Hayakawa) Siver & Wolfe	Hoesan061007G	Hoesan, Illoe-up, Muan-gun, Jeollanam-do, Korea (34°51'37"N, 126°28'58"E)	GU935630	GU935652	KM817941	KM818019	GU935672
<i>M. bronchartiana</i> Compère	Mulyeongari101110J	Mulyeongari, Sumang-ri, Namwon-eup, Seogwipo-si, Jeju-Island, Korea (33°21'52"N,	JQ281517	JQ281511	KM817943	KM818021	JQ281505

			126°41'26"E)					
<i>M. caudata</i> Ivanov emend. Krieger	Bangjukgol062310A	Bangjukgol, Munui-myeon, Cheongwon-gun, Chungcheongbuk-do, Korea (36°32'16"N, 127°29'08"E)	JN991176	JN991185	KM817944	KM818022	JN991167	-
<i>M. corymbosa</i> Asmund & Hiliard	Gungnami072807I	Gungnamji, Buyeo-eup, Buyeo-gun, Chungcheongnam-do, Korea (36°16'12"N, 126°54'46"E)	GU935620	GU935642	KM817947	KM818025	GU935662	-
<i>M. cratis</i> Harris & Bradley	Hoje060207B	Hoje, Daecheong-ri, Cheongha-myeon, Gimje-si, Jeollabuk-do, Korea (35°53'16"N, 126°51'08"E), Mokji, Changnyeong-eup, Changnyeong-gun, Gyeongsangnam-do, Korea (35°21'03"N, 128°27'43"E)	GU935623	GU935645	KM817951	KM818029	GU935665	-
<i>M. elongata</i> Reverdin	Mokjii112407C	Mokjii, Changnyeong-eup, Changnyeong-gun, Gyeongsangnam-do, Korea (35°21'03"N, 128°27'43"E)	GU935621	GU935643	KM817956	KM818034	GU935663	-
<i>M. heterospina</i> Lund	Posan012608A	Posanji, Jindo-eup, Jindo-gun, Jeollanam-do, Korea (34°28'62"N, 126°14'32"E)	GU935617	GU935639	KM817960	KM818038	GU935659	-
<i>M. hexareticulata</i> Jo, Shin, Kim, Siver <i>et</i> Andersen	Mudong072410D	Mudongji, Muji-ri, Hwaseo-myeon, Mungyeongsagi, Gyeongsangbuk-do, Korea (36°36'20"N, 128°18'27"E)	JN991182	JN991191	KM817961	KM818039	JN991173	-
<i>M. insignis</i> Penard	Josanji041710D	Josan, Doam-ri, Dain-myeon, Uiseong-gun, Gyeongsangbuk-do, Korea (36°27'36"N, 128°21'33"E)	JQ955671	JQ955676	KM817963	KM818041	JQ955666	-
<i>M. lacuna</i> Jo, Shin, Kim, Siver <i>et</i>	CCMP2880	Prebersee, Austria (47°11'09"N, 13°51'27"E)	JN991177	JN991186	KM817966	KM818044	JN991168	-

Andersen								
<i>M. mangofera</i> var. <i>foveata</i> (Dürrschmidt) Kristiansen	Hoesan061007I	Hoesan, Illoe-up, Muan-gun, Jeollanam-do, Korea (34°51'37"N, 126°28'58"E)	GU935633	GU935655	KM817970	KM818048	GU935675	-
<i>M. matienkoae</i> Asmund & Kristiansen	Muryeong112807B	Muryeongje, Yeonggwange-up, Yeonggwang-gu, Jeollanam-do, Korea (35°16'59"N, 128°30'56"E),	GU935628	GU935650	KM817971	KM818049	GU935670	-
<i>M. oviformis</i> Nygaard	Jeongdang021410N	Jeongdang, Annyeon-eup, Taean-gun, Chungcheongnam-do, Korea (36°30'32"N, 126°21'43"E)	JN991178	JN991187	KM817976	KM818054	JN991169	-
<i>M. peronoides</i> (Harris) Moneu & Péterfi	Pihyangjeong092609A	Pihyangjeong, Taechang-ri, Taein-myeon, Jeongeup-si, Jeollabuk-do, Korea (35°39'02"N, 126°56'28"E),	JN991180	JN991189	KM817979	KM818057	JN991171	-
<i>M. portiae-ferreae</i> Péterfi & Asmund	Danghal110407B	Dangha, Yongdong-myeon, Iksan-si, Jeollabuk-do, Korea (36°06'19"N, 126°59'36"E)	GU935618	GU935640	KM817980	KM818058	GU935660	-
<i>M. pseudocratis</i> Dürrschmidt	Seungeon120107F	Seungeonji, Seungeon-ri, Anmyeon-eup, Taean-gun, Chungcheongnam-do, Korea (36°31'06"N, 126°20'51"E)	GU935624	GU935646	KM817983	KM818061	GU935666	-
<i>M. pseudomanienkoae</i> Jo, Shin, Kim, Siver <i>et</i> Andersen	Gungnam052507A	Gungnamji, Buyeo-eup, Buyeo-gun, Chungcheongnam-do, Korea (36°16'12"N, 126°54'46"E)	GU935627	GU935649	KM817985	KM818063	GU935669	-
<i>M. punctifera</i> Korshikov	Angumal032010C	Angumal, Deoksan-myeon, Jincheon-gun, 126°54'46"E)	JQ955667	JQ955672	KM817986	KM818064	JQ955662	-

<i>M. sorohexareticulata</i> Jo, Shin, Kim, Gunngnam092709 Siver <i>et</i> Andersen	CCMP1782	Chungcheongbuk-do (36°54'08"N, 127°30'33"E) Gungnamji, Buyeo-eup, Buyeo-gun, Chungcheongnam-do, Korea (36°16'12"N, 128°54'46"E)	JN991183 JN991192 KM817992 KM818070 JN991174 -
<i>M. splendens</i> (West) Playfair	CCMP1782	From freshwater reservoir, Mount Macedon, Victoria Australia (37°28'37"S, 144°33'36"E)	JQ955668 JQ955673 KM817993 KM818071 JQ955663 -
<i>M. stiata</i> var. <i>serrata</i> Harris & Bradley	Hwadong103108B	Hwadongji, Jillye-myeon, Gimhae-si, Gyeongsangnam-do, Korea (35°15'35"N, 128°46'32"E)	GU935622 GU935644 KM817996 KM818074 GU935664 -
<i>M. tonsulata</i> Teiling	Daejej012611J	Daejeji, Jukseong-ri, Singwang-myeon, Zuk-gu, Pohang-si, Gyeongsangbuk-do, Korea (36°08'05"N, 129°15'10"E)	KM817886 KM817921 KM817998 KM818076 KM818112 -
<i>Mallomonas</i> sp.	CCMP476	Salt marsh, England, United Kingdom (54°00'00"N, 04°00'00"W)	JQ281519 JQ251513 - EF432454 -
<i>Synura</i>			
<i>S. americana</i> Kynčlová & Škaloud	Chimu112407C	Chimu, Daesan-myeon, Haman-gun, Gyeongsangnam-do, Korea (35°20'21"N, 128°25'47"E)	KM590551 KM590617 KM590692 KM590769 KM590838 KP268712
<i>S. americana</i> Kynčlová & Škaloud	Chimu112407D	Chimu, Daesan-myeon, Haman-gun, Gyeongsangnam-do, Korea (35°20'21"N,	KM590552 KM590618 KM590693 KM590770 KM590839 KP268713

<i>S. americana</i> Kynčlová & Škaloud	Johae010508F	Johae, Singwang-myeon, Buk-gu, Pohang-si, Gyeongsangbuk-do, Korea (36°06'08"N, 129°15'10"E)	JX455151	JX455155	<b>KM590694</b>	<b>KM590771</b>	JX455147	<b>KP268711</b>
<i>S. americana</i> Kynčlová & Škaloud	A5482	North America	GU325590	-	-	-	GU325180	GU325492
<i>S. americana</i> Kynčlová & Škaloud	CCMP862	Winter's Creek, Keeweenaw County, MI, USA (47°17'30"N, 88°04'19"W)	GU325583	-	-	-	GU325173	GU325485
<i>S. americana</i> Kynčlová & Škaloud	CCMP863	Road ditch near Winter's Creek, Keeweenaw County, MI, USA (47°17'30"N, 88°04'19"W)	GU325584	-	-	-	GU325174	GU325486
<i>S. americana</i> Kynčlová & Škaloud	CCMP865	Winter's Creek, Keeweenaw County, MI, USA (47°17'30"N, 88°04'19"W)	GU325586	-	-	-	GU325176	GU325488
<i>S. americana</i> Kynčlová & Škaloud	Sandgren2	North America	GU325589	-	-	-	GU325179	GU325491
<i>S. americana</i> Kynčlová & Škaloud	UTCC495	Newfoundland Hwy 410, Newfoundland, Canada (49°50'16"N, 56°19'08"W)	GU325595	-	-	-	GU325185	GU325497
<i>S. americana</i> Kynčlová & Škaloud	UTEXLB845	Soil, Bloomington, IN, USA	GU325592	-	-	-	GU325182	GU325494
<i>S. asmundiae</i> (Cronberg & Kristiansen) Škaloud, Kristiansen & Škaloudová	S90D10	Rutajärvi, Finland (61°54'50"N, 26°04'52"E)	<b>KM590553</b>	<b>KM590619</b>	<b>KM590695</b>	-	<b>KM590840</b>	<b>KP268729</b>
<i>S. asmundiae</i> (Cronberg & Kristiansen) Škaloud, Kristiansen & Škaloudová	S90D11	Rutajärvi, Finland (61°54'50"N, 26°04'52"E)	HF549069	-	-	-	HF549079	<b>KP268730</b>

<i>S. hjoerki</i> (Cronberg & Kristiansen) Škaloud, Kristiansen & Škaloudová	SC57A6	A small canal, Store Mosse, Sweden (57°18'23"N, 14°00'45"E)	HF549070	-	-	-	-	HF549080 KP268731
<i>S. conopea</i> Kynčlová & Škaloud	Gonggeomji2103108C	Gonggeom, Yangseong-ri, Gonggeom-myeon, Sangju-si, Gyeongsangbuk-do, Korea (36°29'42"N, 128°07'29"E)	KM590554	KM590620	KM590696	KM590772	KM590841	KP268693
<i>S. conopea</i> Kynčlová & Škaloud	Gonggeomji4031909C	Gonggeom, Yangseong-ri, Gonggeom-myeon, Sangju-si, Gyeongsangbuk-do, Korea (36°29'42"N, 128°07'29"E)	KM590555	KM590621	KM590697	KM590773	KM590842	KP268692
<i>S. conopea</i> Kynčlová & Škaloud	Josan041808A	Josan, Doam-ri, Dain-myeon, Uiseong-gun, Gyeongsangbuk-do, Korea (36°27'36"N, 128°21'33"E)	KM590556	KM590622	KM590698	KM590774	KM590843	KP268691
<i>S. conopea</i> Kynčlová & Škaloud	Sugyeji041808B	Sugye, Dain-myeon, Uiseong-gun, Gyeongsangbuk-do, Korea (36°24'73"N, 128°25'13"E)	KM590557	KM590623	KM590699	KM590775	KM590844	KP268690
<i>S. conopea</i> Kynčlová & Škaloud	Yeonseong120807E	Yeonseong, Yeonseong-ri, Yeonggwang-eup, Yeonggwang -gun, Jeollanam-do, Korea (35°17'16"N, 126°31'01"E)	KM590558	KM590624	KM590700	KM590776	KM590845	KP268689
<i>S. conopea</i> Kynčlová & Škaloud	CCMP859	Southeast bridge pond, Washington County, AR, USA	GU325580	-	-	-	GU325170	GU325482 GU338121
<i>S. conopea</i> Kynčlová & Škaloud	KNU13	Dalseonggyo, Gyeongju-si, Gyeongsangbuk-do, Korea (36°03'07"N, 129°14'17"E)	GU325529	-	-	-	GU325116	GU325430 GU338069

<i>S. conopea</i> Kynčlová & Škaloud	NIES1007		Tomakomai, Hokkaido, Japan (42°39'58"N, 141°36'43"E)	GU325578	-	-	GU325168	GU325479	GU338119
<i>S. curispina</i> (Petersen & Hansen)	Gaekmang111107A	Asmund	Gaekmang, Udeok-ri, Deokcheon-myeon, Jeongeup-si, Jeollabuk-do, Korea (35°37'34"N, 128°50'41"E)	KM590559	KM590625	KM590701	KM590777	KM590846	KP268734
<i>S. curispina</i> (Petersen & Hansen)	Giududeul090311A	Asmund	Giududeul, Sowon-myeon, Taean-gun, Chungcheongnam-do, Korea (37°32'41"N, 127°05'16"E)	KM590560	KM590626	KM590702	KM590778	KM590847	KP268736
<i>S. curispina</i> (Petersen & Hansen)	Guungnam052507A	Asmund	Guungnamji, Buyeo-eup, Buyeo-gun, Chungcheongnam-do, Korea (36°16'12"N, 126°54'46"E)	KM590561	KM590627	KM590703	KM590779	KM590848	KP268735
<i>S. curispina</i> (Petersen & Hansen)	Hanjeong080611B	Asmund	Hanjeong, Girin-ri, Soseong-myeon, Jeongeup-si, Jeollabuk-do, Korea (35°33'55"N, 126°46'30"E)	KM590562	KM590628	KM590704	KM590780	KM590849	KP268733
<i>S. curispina</i> (Petersen & Hansen)	SAG2992	Asmund	Berlin-Zehlendorf, Nikolassee, Germany	GU325515	-	-	GU325102	GU325415	GU338151
<i>S. echinulata</i> Korshikov	CCMP853	Asmund	Bog near calumet, MT, USA (47°14'38"N, 88°31'12"W)	KM590563	KM590629	KM590705	KM590781	KM590850	KP268754
<i>S. echinulata</i> Korshikov	SAG1592	S. glabra	Berlin-Spandau, Swamp in forest, Germany (52°27'25"N, 13°16'43"E)	GU325513	-	-	GU325101	GU325414	GU338154
<i>S. glabra</i> Korshikov emend.	Bonggye101407K	Kynčlová & Škaloud	Bonggye, Bonggye-ri, Dudong-myeon, Ulju-gun, Ulsan, Korea (35°40'55"N, 129°12'43"E)	KM590564	KM590630	KM590706	KM590782	KM590851	KP268722

<i>S. glabra</i> Korshikov emend. Kynčlová & Škaloud	Cheonma041908B	Cheonma, Yugok-ri, Yuga-myeon, Dalseong-gun, Gyeongsangbuk-do, Korea (36°54'29"N, 128°46'17"E)	KM590565 KM590631 KM590707 KM590783 KM590852 KP268716
<i>S. glabra</i> Korshikov emend. Kynčlová & Škaloud	Dohak11107C	Dohak, Udeok-ri, Deokcheon-myeon, Jeongeup- si, Jeollabuk-do, Korea (35°37'21"N, 128°49'37"E)	KM590566 KM590632 KM590709 KM590785 KM590853 KP268721
<i>S. glabra</i> Korshikov emend. Kynčlová & Škaloud	Dohak11107E	Dohak, Udeok-ri, Deokcheon-myeon, Jeongeup- si, Jeollabuk-do, Korea (35°37'21"N, 128°49'37"E)	KM590566 KM590632 KM590709 KM590785 KM590853 KP268725
<i>S. glabra</i> Korshikov emend. Kynčlová & Škaloud	Eunseon11107C	Eunseon, Eunseon-ri, Yeongwon-myeon, Jeongeup-si, Jeollabuk-do, Korea (35°37'21"N, 126°49'37"E)	KM590567 KM590633 KM590710 KM590786 KM590854 KP268720
<i>S. glabra</i> Korshikov emend. Kynčlová & Škaloud	Geumma020610B	Geumma, Geumma-myeon, Iksan-si, Jeollabuk- do, Korea (36°00'02"N, 127°03'09"E)	KM590568 KM590634 KM590711 KM590787 KM590855 KP268718
<i>S. glabra</i> Korshikov emend. Kynčlová & Škaloud	Gungnam3102807A	Gungnamji, Buyeo-eup, Buyeo-gun, Chungcheongnam-do, Korea (36°16'12"N, 126°54'46"E)	KM590569 KM590635 KM590712 KM590788 KM590856 KP268727
<i>S. glabra</i> Korshikov emend. Kynčlová & Škaloud	Hoesan061007L	Hoesan, Illoe-up, Muan-gun, Jeollanam-do, Korea (34°51'37"N, 126°28'58"E)	KM590570 KM590636 KM590713 KM590789 KM590857 KP268719
<i>S. glabra</i> Korshikov emend. Kynčlová & Škaloud	Hwangsang012508A	Hwangsang, Haeje-myeon, Muan-gun, Jeollanam- do, Korea (35°06'02"N, 126°19'30"E)	KM590571 KM590637 KM590714 KM590790 KM590858 KP268724

<i>S. glabra</i> Korshikov emend. Kynčlová & Škaloud	Jakeunmeok052407B	Jakeunmeok, Seungeon-ri, Annyeon-eup, Taean-gun, Chungcheongnam-do, Korea (36°35'30"N, 126°40'34"E)	KM590572	KM590638	KM590715	KM590791	KM590859	KP268728
<i>S. glabra</i> Korshikov emend. Kynčlová & Škaloud	Oryong020610B	Oryong, Seogodo-ri, Geumma-myeon, Iksan-si, Jeollabuk-do, Korea (35°56'11"N, 127°03'56"E)	KM590573	KM590639	KM590716	KM590792	KM590860	KP268717
<i>S. glabra</i> Korshikov emend. Kynčlová & Škaloud	Seoseong060307E	Seoseong, Deokgi-dong, Iksan-si, Jeollabuk-do, Korea (35°58'50"N, 127°02'23"E)	KM590574	KM590640	KM590717	KM590793	KM590861	KP268723
<i>S. glabra</i> Korshikov emend. Kynčlová & Škaloud	Yeondong060207B	Yeondong, Geumma-myeon, Iksan-si, Jeollabuk-do, Korea (35°59'04"N, 127°02'43"E)	KM590575	KM590641	KM590718	KM590794	KM590862	KP268726
<i>S. glabra</i> Korshikov emend. Kynčlová & Škaloud	Yukgeum110407B	Yukgeum, Jenam-ri, Yeosan-myeon, Iksan-si, Jeollabuk-do, Korea (36°03'26"N, 127°04'21"E)	KM590576	KM590642	KM590719	KM590795	KM590863	KP268715
<i>S. lanceolata</i> Jo, Kim, Siver & Shin sp. nov.	Mangaeji040409A	Mangaeji, Bibong-myeon, Cheongyang-gun, Chungcheongnam-do, Korea (36°33'35"N, 126°45'42"E)	KM590577	KM590643	KM590720	KM590796	KM590864	KP268700
<i>S. lanceolata</i> Jo, Kim, Siver & Shin sp. nov.	Nojeok020610A	Nojeok, Chumpo-myeon, Iksan-si, Jeollabuk-do, Korea (35°56'48"N, 127°01'43"E)	KM590578	KM590644	KM590721	KM590797	KM590865	KP268701
<i>S. lanceolata</i> Jo, Kim, Siver & Shin sp. nov.	Oksan120807A	Oksan, Oksan-ri, Mujang-myeon, Gochang-gun, Jeollabuk-do, Korea (35°24'05"N, 126°35'19"E)	KM590579	KM590645	KM590722	KM590798	KM590866	KP268699
<i>S. lanceolata</i> Jo & Shin	CNU49	Ocksan pond, Gochang-gun, Jeollabuk-do, Korea	GU325575	-	-	-	GU325164	GU325477
<i>S. longitubularis</i> Jo, Kim, Siver & Shin sp. nov.	Jeongsan070607A	Jeongsan, Jeongsan-myeon, Cheongyang-gun, Chungcheongnam-do, Korea (36°25'13"N,	KM590580	KM590646	KM590723	KM590799	KM590867	KP268740

<i>S. longitubularis</i> Jo, Kim, Siver & Shin <i>sp. nov.</i>	Sinsan101709B	Sinsan, Dain-myeon, Uiseong-gun, Gyeongsangbuk-do, Korea (36°25'24"N, 128°22'52"E),	<b>KM590581</b>	<b>KM590647</b>	<b>KM590724</b>	<b>KM590800</b>	<b>KM590868</b>	<b>KP268739</b>
<i>S. macracantha</i> (Petersen & Hansen) S90B5 Asmund	S71B2	Tehriselkä, Finland (64°45'23"N, 26°29'08"E)	HF549064	<b>KM590648</b>	<b>KM590725</b>	-	HF549075	<b>KP268732</b>
<i>S. macropora</i>	Santaek072410C	Santaek, Santaek-ri, Yonggung-myeon, Yecheon- gun, Gyeongsangbuk-do, Korea (36°36'32"N, 128°17'55"E)	<b>KM590582</b>	-	<b>KM590726</b>	-	<b>KM590869</b>	<b>KP268714</b>
<i>S. mammillosa</i> Takahashi	S18	Dongbaek-garden, Seonheul-ri, Jocheon-eup, Jeju-si, Jeju-Island, Korea (33°30'31"N, 126°43'03"E)	<b>KM590584</b>	<b>KM590650</b>	<b>KM590728</b>	<b>KM590802</b>	<b>KM590871</b>	<b>KP268748</b>
<i>S. mammillosa</i> Takahashi	S19	Dongbaek-garden, Seonheul-ri, Jocheon-eup, Jeju-si, Jeju-Island, Korea (33°30'31"N, 126°43'03"E)	<b>KM590585</b>	<b>KM590651</b>	<b>KM590729</b>	<b>KM590803</b>	<b>KM590872</b>	<b>KP268749</b>
<i>S. mammillosa</i> Takahashi	S89C3	Unnamed lake near Koivula, Finland (62°15'01"N, 26°34'48"E)	HF549066	<b>KM590652</b>	-	-	-	<b>KP268752</b>
<i>S. mammillosa</i> Takahashi	S96B5	Loch Braigh nan Allt, Scotland , United Kingdom (58°00'59"N, 05°04'48"W)	-	<b>KM590653</b>	<b>KM590730</b>	-	-	<b>KP268751</b>
<i>S. mammillosa</i> Takahashi	SIE105A	Caha lakes, Beara, Ireland (51°43'18"N, 10	HF549065	<b>KM590654</b>	-	-	HF549076	<b>KP268753</b>

		09°39'37" W)
<i>S. molispina</i> (Petersen & Hansen)	S71C10	Podhradská pool, Czech Republic (50°27'40"N, 14°54'42"E)
Péterfi & Momeu		
<i>S. multidentata</i> (Balonov & Kuzmin)	S90C11	Heinäjärvi, Finland (62°52'52"N, 25°29'50"E)
Péterfi & Momeu		
<i>S. petersenii</i> Korshikov emend.	Buje100307A	Buje, Gapje-dong, Gyeongsan-si, Gyeongsangbuk-do, Korea (35°49'54"N, 128°46'14"E)
Škaloud & Kynčlová		
<i>S. petersenii</i> Korshikov emend.	Gangok111107C	Gangok, Gangok-myeon, Jeongeup-si, Jeollabuk-do, Korea (35°43'08"N, 126°57'45"E)
Škaloud & Kynčlová		
<i>S. petersenii</i> Korshikov emend.	Saeraeul103109B	Saerae, Haenggu-dong, Wonju-si, Gangwon-do, Korea (37°20'34"N, 128°00'23"E)
Škaloud & Kynčlová		
<i>S. petersenii</i> Korshikov emend.	Swaeji103109I	Swaeji, Haenggu-dong, Wonju-si, Gangwon-do, Korea (37°21'09"N, 128°00'14"E)
Škaloud & Kynčlová		
<i>S. petersenii</i> Korshikov emend.	Yongseong112407A	Yongseong, Yongseong-ri, Chilseo-myeon, Haman-gun, Gyeongsangnam-do, Korea (35°20'03"N, 128°29'54"E)
Škaloud & Kynčlová		
<i>S. petersenii</i> Korshikov emend.	Youngji101407A	Youngji, Gyeongju-si, Gyeongsangbuk-do, Korea (35°45'02"N, 129°17'52"E)
Škaloud & Kynčlová		
<i>S. petersenii</i> Korshikov emend.	Youngji101407B	Youngji, Gyeongju-si, Gyeongsangbuk-do, Korea (35°45'02"N, 129°17'52"E)
Škaloud & Kynčlová		

<i>S. petersonii</i> Korshikov emend. Škaloud & Kynčlová	CNU39	Yongseong, Yongseong-ri, Chilseo-myeon, Haman-gun, Gyeongsangnam-do, Korea (35°20'03"N, 128°29'54"E)	GU325565	-	-	GU325154	GU325467	GU338106
<i>S. soroconeopea</i> Jo, Kim, Siver & Shin <i>sp. nov.</i>	Jakeun030207B	Jakeunmeok, Seungeon-ri, Anmyeon-eup, Taeang- gun, Chungcheongnam-do, Korea (36°35'30"N, 128°40'34"E)	KM590592	KM590663	KM590739	KM590811	KM590879	KP268686
<i>S. soroconeopea</i> Jo, Kim, Siver & Shin <i>sp. nov.</i>	Mudeomsil102007A	Mudeom, Gaejeon-ri, Hoengseong-eup, Hoengseong-gun, Gangwon-do, Korea (37°30'17"N, 128°01'19"E)	KM590593	KM590664	KM590740	KM590812	KM590880	KP268682
<i>S. soroconeopea</i> Jo, Kim, Siver & Shin <i>sp. nov.</i>	Sorigae122107A	Sori, Cheongyang-eup, Cheongyangang-guun, Chungcheongnam-do, Korea (36°27'55"N, 126°46'33"E)	KM590594	KM590665	KM590741	KM590813	KM590881	KP268687
<i>S. soroconeopea</i> Jo, Kim, Siver & Shin <i>sp. nov.</i>	Sorigae122107C	Sori, Cheongyang-eup, Cheongyangang-guun, Chungcheongnam-do, Korea (36°27'55"N, 126°46'33"E)	KM590595	KM590666	KM590742	KM590814	KM590882	KP268688
<i>S. soroconeopea</i> Jo, Kim, Siver & Shin <i>sp. nov.</i>	Sotan012410A	Sotan, Geumsseong-myeon, Geumsan-guun, Chungcheongnam-do, Korea (36°07'54"N, 126°46'33"E)	KM590596	KM590667	KM590743	KM590815	KM590883	KP268685
<i>S. soroconeopea</i> Jo, Kim, Siver & Shin <i>sp. nov.</i>	Sugyeji041808J	Sugye, Dain-myeon, Uiseong-guun, Gyeongsangbuk-do, Korea (36°24'73"N, 128°25'13"E)	KM590597	KM590668	KM590744	KM590816	KM590884	KP268684

<i>S. soroconepea</i> Jo, Kim, Siver & Shin <i>sp. nov.</i>	Yukgeum110407A	Yukgeum, Jenam-ri, Yeosan-myeon, Iksan-si, Jeollabuk-do, Korea (36°03'26"N, 127°04'21"E)	<b>KM590598</b>	<b>KM590669</b>	<b>KM590745</b>	<b>KM590817</b>	<b>KM590835</b>	<b>KP268683</b>
<i>S. soroconepea</i> Jo, Kim, Siver & Shin <i>sp. nov.</i>	CNU01	Samhwa pond, Gwangju-si, Korea	GU325530	-	-	GU325117	GU325431	GU338070
<i>S. soroconepea</i> Jo, Kim, Siver & Shin <i>sp. nov.</i>	CNU10	Bihaksan, Singwang-myeon, Buk-gu, Pohang-si, Gyeongsangbuk-do, Korea	GU325538	-	-	GU325125	GU325439	GU338078
<i>S. sphagnicola</i> (Korshikov) Korshikov	CCMP1705	Keeweenaw County, MI, USA (47°17'30"N, 88°04'19"W)	U73221	-	-	-	-	EF165197
<i>S. sphagnicola</i> (Korshikov) Korshikov	CCMP2897	Loibersbacher Teich 2, Austria (47°44'11"N, 13°18'02"E)	-	-	-	GU325100	GU325413	GU338153
<i>S. sphagnicola</i> (Korshikov) Korshikov	JYS001	Unknown	DQ980485	DQ980475	-	-	-	-
<i>S. sphagnicola</i> (Korshikov) Korshikov	Angol1129J2	Angol, Jeongsan-myeon, Cheongyang-gun, Chungcheongnam-do, Korea (36°25'13"N, 126°56'16"E)	KT364487	KT364489	KT364491	KT364493	KT364495	KT364497
<i>S. sphagnicola</i> (Korshikov) Korshikov	Hudong1129J2	Hudong, Jeongsan-myeon, Cheongyang-gun, Chungcheongnam-do, Korea (36°25'31"N, 126°56'26"E)	KT364488	KT364490	KT364492	KT364494	KT364496	KT364498
<i>S. spinosa</i> Korshikov Korshikov	Anha103010C	Anha, Jisa-myeon, Imsil-gun Jeollabuk-do, Korea (35°34'52"N, 127°21'13"E)	<b>KM590599</b>	<b>KM590670</b>	<b>KM590746</b>	<b>KM590818</b>	<b>KM590836</b>	<b>KP268744</b>
<i>S. spinosa</i> Korshikov	Hakgok103109B	Hakgok, Ucheon-myeon, Hoengseong -gun,	<b>KM590600</b>	<b>KM590671</b>	<b>KM590747</b>	<b>KM590819</b>	<b>KM590887</b>	<b>KP268743</b>

<i>S. spinosa</i> Korshikov	Osan103010A	Osan, Sanseo-myeon, Jangsu-gun, Jeollabuk-do, Korea (35°35'03"N, 127°24'34"E)	KM590601	KM590672	KM590748	KM590820	KM590888	KP268745
<i>S. spinosa</i> Korshikov	Sandoji011109C	Sandoji, Hwangsang-dong, Gimje-si, Jeollabuk-do, Korea (35°46'21"N, 126°55'54"E)	KM590602	KM590673	KM590749	KM590821	KM590889	KP268742
<i>S. spinosa</i> Korshikov	S74D2	Pískovny kanál, Veselské sand quarries, Czech Republic (49°09'13"N, 14°42'35"E)	-	-	-	-	HF549081	KP268746
<i>S. splendida</i> Korshikov	S90E4	Rutajärvi, Finland (61°54'50"N, 26°04'53"E)	HF549071	-	-	-	HF549082	KP268761
<i>S. splendida</i> Korshikov	S90E6	Rutajärvi, Finland (61°54'50"N, 26°04'53"E)	KM590603	KM590674	KM590750	-	KM590890	KP268762
<i>S. sungminbooi</i> Jo & Shin	Buyeom032010A	Buyeon, Buyeon-ri, Bugi-myeon, Cheongwon-gun, Chungcheongbuk-do, Korea (36°44'12"N, 127°34'18"E)	KM590604	KM590675	KM590751	KM590822	KM590891	KP268696
<i>S. sungminbooi</i> Jo, Kim, Siver & Shin <i>sp. nov.</i>	Geomuyeok040507	Geomu, Geomuyeok-ri, Byeonggok-myeon, Yeongdeok-gun, Gyeongsangbuk-do, Korea (36°35'18"N, 129°23'35"E)	KM590605	KM590676	KM590752	KM590823	KM590892	KP268697
<i>S. sungminbooi</i> Jo, Kim, Siver & Shin <i>sp. nov.</i>	Geumgok020610A	Geumgok, Duyeo-ri, Yeosan-myeon, Iksan-si, Jeollabuk-do, Korea (36°04'10"N, 127°05'02"E)	KM590606	KM590677	KM590753	KM590824	KM590893	KP268698
<i>S. sungminbooi</i> Jo, Kim, Siver & Shin <i>sp. nov.</i>	Seokjangji032010A	Seokjang, Seokjang-ri, Deoksan-myeon, Jincheon-gun, Chungcheongbuk-do, Korea (36°53'37"N, 127°31'36"E)	KM590607	KM590678	KM590754	KM590825	KM590894	KP268695
<i>S. sungminbooi</i> Jo, Kim, Siver & Shin	Seongdong120807A	Seongdong, Yeonseong-ri, Yeonggwange-up, Seongdong, Yeonseong-ri, Yeonggwange-up,	KM590608	KM590679	KM590755	KM590826	KM590895	KP268694

<i>sp. nov.</i>	Yeonggwang-gun, Jeollanam-do, Korea (35°16'52"N, 126°31'42"E)						
<i>S. sunghminbooi</i> Jo, Kim, Siver & Shin <i>sp. nov.</i>	CNU12	Geomu, Geomuyeok-ri, Byeonggok-myeon, Yeongdeok-gun, Gyeongsangbuk-do, Korea (36°35'18"N, 129°23'35"E)	GU325540	-	-	GU325127	GU325441 GU338080
<i>S. sunghminbooi</i> Jo, Kim, Siver & Shin <i>sp. nov.</i>	CNU47	Seongdong, Yeonseong-ri, Yeonggwange-up, Yeonggwang-gun, Jeollanam-do, Korea (35°16'52"N, 126°31'42"E)	GU325573	-	-	GU325162	GU325475 GU338114
<i>S. truttae</i> (Siver) Škaloud & Kynčlová	Hanjeong080611J	Hanjeong, Girin-ri, Soseong-myeon, Jeongeup-si, Jeollabuk-do, Korea (35°33'55"N, 126°46'30"E)	KM590609	KM590680	KM590756	KM590827	KM590896 KP268702
<i>S. truttae</i> (Siver) Škaloud & Kynčlová	Jangjuk032611J	Jangjuk, Hyanghan-ri, Eomsa-myeon, Gyeryong- si, Chungcheongnam-do, Korea (36°16'31"N, 127°12'38"E)	KM590610	KM590681	KM590757	KM590828	KM590897 KP268703
<i>S. truttae</i> (Siver) Škaloud & Kynčlová	Nemcova 2	Czech Republic	GU325598	-	KM590758	GU325188	GU325500 GU338138
<i>S. truttae</i> (Siver) Škaloud & Kynčlová	Nemcova 5	Czech Republic	GU325600	-	-	GU325190	GU325502 GU338140
<i>S. truttae</i> (Siver) Škaloud & Kynčlová	Nemcova 6	Czech Republic	GU325601	-	-	GU325191	GU325503 GU338141
<i>S. uvella</i> Ehrenberg emend. Korshikov	Gahang033107A	Gahang, Gahang-ri, Yueo-myeon, Changnyeong- gun, Gyeongsangnam-do, Korea (35°31'27"N,	KM590611	KM590682	KM590759	KM590829	KM590898 KP268756

<i>S. uvella</i> Ehrenberg emend. Korshikov	Geumsan032709D	Geumsan, Gongsan-myeon, Naju-si, Jeollanam-do, Korea (34°57'20"N, 126°35'25"E)	<b>KM590612</b>	KM590683	<b>KM590760</b>	KM590830	KM590899	<b>KP268757</b>
<i>S. uvella</i> Ehrenberg emend. Korshikov	Gonggeom2103108A	Gonggeom, Yangseong-ri, Gonggeom-myeon, Sangu-si, Gyeongsangbuk-do, Korea (36°29'42"N, 128°07'29"E)	<b>KM590613</b>	KM590684	<b>KM590761</b>	KM590831	KM590900	<b>KP268758</b>
<i>S. uvella</i> Ehrenberg emend. Korshikov	AI3.377	Lake of the Clouds, Ontonagon, MI , USA (46°48'09"N, 89°45'08"W)	<b>KM590614</b>	KM590685	-	-	<b>KM590901</b>	<b>KP268759</b>
<i>S. uvella</i> Ehrenberg emend. Korshikov	CCMP870	Volo Bog Nature Preserve, McHenry County, IL, USA (42°21'05"N, 88°11'04"W)	<b>KM590615</b>	KM590686	-	-	AF015586	<b>KP268760</b>
<i>S. uvella</i> Ehrenberg emend. Korshikov	CCMP871	Tobacco River, Keeweenaw County, MI, USA (43°49'00"N, 84°23'18"W)	U73222	-	-	-	EF165192	-
<i>S. uvella</i> Ehrenberg emend. Korshikov	CNU53	Daepyeong swamp, Beopsu-myeon, Haman-gun, Gyeongsangnam-do, Korea (35°20'23"N, 128°20'10"E)	GU325514	-	-	GU325103	GU325416	GU338056
<i>Synura</i> sp.	450	Unknown	-	<b>KM590687</b>	-	-	<b>KM590902</b>	<b>KP268741</b>
<i>Synura</i> sp.	CCAC0052	Cologne, Germany	GU325606	-	-	GU325196	GU325508	GU338146
<i>Synura</i> sp.	CCMP847	Winter's Creek, Keeweenaw County, MI, USA (47°17'30"N, 88°04'19"W)	EF165128	-	-	-	EF165196	-
<i>Synura</i> sp.	CCMP869	Tobacco River, Keeweenaw County, MI, USA (43°49'00"N, 84°23'18"W)	GU325587	<b>KM590688</b>	<b>KM590762</b>	GU325177	GU325489	GU338128

<i>Synura</i> sp.	CCMP2898	Wallersee, Austria (47°54'28"N, 13°10'06"E)	GU325596	-	-	GU325186	GU325498	GU338136
<i>Synura</i> sp.	Gungnam1072807A	Gungnamji, Buyeo-eup, Buyeo-gun, Chungcheongnam-do, Korea (36°16'12"N, 126°54'46"E)	<b>KM590616</b>	<b>KM590689</b>	<b>KM590763</b>	<b>KM590832</b>	<b>KM590903</b>	<b>KP268737</b>
<i>Synura</i> sp.	Igyo111107A	Igyo, Jeongeup-si, Jeollabuk-do, Korea (35°41'14"N, 126°54'18"E)	JX455148	JX455152	<b>KM590764</b>	<b>KM590833</b>	JX455144	<b>KP268738</b>
<i>Synura</i> sp.	UTEXLB239	Unknown	GU325591	-	-	GU325181	GU325493	GU338135
<i>Synura</i> sp.	WA18K_U	Austria	GU325597	-	-	GU325187	GU325499	GU338137
<i>Tessellaria</i>								
<i>Tessellaria laponica</i> Skuja (Škaloud, Christiansen & Škaloudová)	S59C4	Helgassjön, Sweden (56°57'22"N, 14°42'58"E)	HF549063	<b>KM590690</b>	-	-	HF549074	<b>KP268681</b>
<i>Tessellaria volvocina</i> Playfair	CCMP1781	Freshwater reservoir, Mount Macedon, Victoria, Australia (37°28'37"S, 144°33'36"E)	EF165119	<b>KM590691</b>	<b>KM590765</b>	<b>KM590834</b>	EF165199	<b>KP268680</b>
<b>Outgroup</b>								
<i>Chromulina</i> sp.	SAG 17.97	Bog pool, Oberturgl/Oetztal, Tyrol, Austria (46°52'12"N, 11°01'38"E)	EF165103	GU935638	<b>KM590766</b>	<b>KM590835</b>	EF165151	-
<i>Ochromonas danica</i> Pringsheim	SAG 933.7	Bog pool, Everdrup, Denmark (55°12'12"N, 11°56'47"E)	JQ281514	GU935636	<b>KM590767</b>	<b>KM590836</b>	GU935657	-
<i>Ochromonas</i> sp.	SAG 933.10	Pond, Solling, Germany (51°43'22"N, 09°37'18"E)	EF165109	GU935637	<b>KM590768</b>	<b>KM590837</b>	GU935658	-

New sequences are indicated in bold type.